



Scientific Excellence • Resource Protection & Conservation • Benefits for Canadians
Excellence scientifique • Protection et conservation des ressources • Bénéfices aux Canadiens

**Rockfish Sampling Cruise:
F/V OCEAN SELECTOR, July 12
to 30, 1989**

G. E. Gillespie and B. M. Leaman

Department of Fisheries and Oceans
Biological Sciences Branch
Pacific Biological Station
Nanaimo, British Columbia V9R 5K6

February 1990

**Canadian Manuscript Report of
Fisheries and Aquatic Sciences
No. 2061**



Fisheries
and Oceans

Pêches
et Océans

Canada

Canadian Manuscript Report of Fisheries and Aquatic Sciences

Manuscript reports contain scientific and technical information that contributes to existing knowledge but which deals with national or regional problems. Distribution is restricted to institutions or individuals located in particular regions of Canada. However, no restriction is placed on subject matter, and the series reflects the broad interests and policies of the Department of Fisheries and Oceans, namely, fisheries and aquatic sciences.

Manuscript reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report is abstracted in *Aquatic Sciences and Fisheries Abstracts* and indexed in the Department's annual index to scientific and technical publications.

Numbers 1-900 in this series were issued as Manuscript Reports (Biological Series) of the Biological Board of Canada, and subsequent to 1937 when the name of the Board was changed by Act of Parliament, as Manuscript Reports (Biological Series) of the Fisheries Research Board of Canada. Numbers 901-1425 were issued as Manuscript Reports of the Fisheries Research Board of Canada. Numbers 1426-1550 were issued as Department of Fisheries and the Environment, Fisheries and Marine Service Manuscript Reports. The current series name was changed with report number 1551.

Manuscript reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page. Out-of-stock reports will be supplied for a fee by commercial agents.

Rapport manuscrit canadien des sciences halieutiques et aquatiques

Les rapports manuscrits contiennent des renseignements scientifiques et techniques qui constituent une contribution aux connaissances actuelles, mais qui traitent de problèmes nationaux ou régionaux. La distribution en est limitée aux organismes et aux personnes de régions particulières du Canada. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques du ministère des Pêches et des Océans, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports manuscrits peuvent être cités comme des publications complètes. Le titre exact paraît au-dessus du résumé de chaque rapport. Les rapports manuscrits sont résumés dans la revue *Résumés des sciences aquatiques et halieutiques*, et ils sont classés dans l'index annuel des publications scientifiques et techniques du Ministère.

Les numéros 1 à 900 de cette série ont été publiés à titre de manuscrits (série biologique) de l'Office de biologie du Canada, et après le changement de la désignation de cet organisme par décret du Parlement, en 1937, ont été classés comme manuscrits (série biologique) de l'Office des recherches sur les pêcheries du Canada. Les numéros 901 à 1425 ont été publiés à titre de rapports manuscrits de l'Office des recherches sur les pêcheries du Canada. Les numéros 1426 à 1550 sont parus à titre de rapports manuscrits du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 1551.

Les rapports manuscrits sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre. Les rapports épuisés seront fournis contre rétribution par des agents commerciaux.

91-00877/

Canadian Manuscript Report of
Fisheries and Aquatic Sciences No. 2061

February 1990

ROCKFISH SAMPLING CRUISE:
F/V OCEAN SELECTOR, JULY 12 TO 30, 1989

by

G. E. Gillespie and B. M. Leaman

Department of Fisheries and Oceans
Biological Sciences Branch
Pacific Biological Station
Nanaimo, British Columbia V9R 5K6

(c) Minister of Supply and Services Canada 1990

Cat. No. Fs 97-4/2061E

ISSN 0706-6473

Correct citation for this publication:

Gillespie, G. E. and B. M. Leaman. 1990. Rockfish sampling
cruise: F/V OCEAN SELECTOR, July 12 to 30, 1989. Can. MS
Rep. Fish. Aquat. Sci. 2061: 89 p.

ABSTRACT

Gillespie, G. E. and B. M. Leaman. 1990. Rockfish sampling cruise: F/V OCEAN SELECTOR, July 12 to 30, 1989. Can. MS Rep. Fish. Aquat. Sci. 2061: 89 p.

Preliminary results from a research cruise to collect biological samples of commercially important rockfishes in British Columbia waters are presented. Decreased trip limits in the commercial fishery have limited opportunities to obtain discrete samples from individual management areas. This cruise was designed to provide the population data for rockfishes, primarily Pacific ocean perch (Sebastes alutus).


A total of 6,775 Pacific ocean perch and 1,082 other rockfish were sampled from traditional fishing grounds off the west coasts of the Queen Charlotte Islands and Vancouver Island, and in Queen Charlotte Sound.

RÉSUMÉ

Gillespie, G. E. and B. M. Leaman. 1990. Rockfish sampling cruise: F/V OCEAN SELECTOR, July 12 to 30, 1989. Can. MS Rep. Fish. Aquat. Sci. 2061: 89 p.

Les résultats préliminaires d'une croisière de recherche ayant pour but de faire des prélèvements biologiques des rascasses ayant une importance commerciale dans les eaux de la Colombie-Britannique sont présentés. La réduction des limites de prises par sortie dans la pêche commerciale a restreint les occasions d'obtenir des échantillons discrets de chacun des secteurs d'exploitation. Cette croisière visait à réunir des données sur les populations de rascasses, et surtout sur la sébaste du Pacifique (Sebastes alutus).

On a échantillonné 6 775 sébastes du Pacifique et 1 082 autres rascasses dans les champs de pêche traditionnels au large de la côte ouest des Iles de la Reine-Charlotte et de l'île Vancouver, ainsi que dans le détroit de la Reine-Charlotte.



INTRODUCTION

The Pacific Region Groundfish Management Plan (Anon. 1989) attempts to balance the competing needs of stock conservation and economic return for the B.C. groundfish industry. One of the goals of the Plan is to have the fishery occupy as much of the year as possible. The mechanisms chosen to accomplish this are individual vessel trip limits and quarterly quotas, scaled to vessel size, time of year and species. In general, the Plan has helped to prevent large overruns in quotas but it has also restricted sampling opportunities, as trip limits are reduced in parts of the year. These lower limits (2-25 t) mean that vessels must combine many such limits, by species and areas, to provide landings similar in total quantity to those prior to reduction (100-150 t). Combination of small catches from different species and areas in vessel tanks or holds means that discrete samples from management units become increasingly more difficult to obtain. Since one explicit management goal is to extend the fishery for as long as possible throughout the year, the mechanisms of management act to decrease the availability of information upon which to base that management. This is an inevitable result of complex stock structures, variable productivity levels, and management goals. An alternative would be to have open fisheries until quotas are fully subscribed. However, there would be obvious biases associated with particular vessel classes and times of year.

To overcome this difficulty, the Offshore Rockfish Program has initiated a series of synoptic sampling cruises, to obtain biological samples from shelf and slope rockfish stocks. Timing of the cruises depends on sampling goals, but for population data we attempt to sample when males and females are in approximately equal sex ratio on the fishing grounds (June-August). Samples taken at other times of year are subject to bias as a result of seasonal differences in availability and movement of the sexes (Leaman 1985).

This report presents some results of the slope rockfish sampling cruise conducted in July, 1989. Age-specific results will be contained in future reports.

MATERIALS AND METHODS

Vessel and gear

Samples were collected aboard the F/V OCEAN SELECTOR, a 47.9 m Canadian commercial stern trawler equipped for both bottom

and midwater fishing (Appendix Table 1, Appendix Figs. 1 and 2).

Sampling methods

Sampling stations were selected by the vessel master, based on his knowledge of the area, for designated survey areas between Dixon Entrance and southwest Vancouver Island (Figs. 1-3). We attempted to obtain two samples for each 37 m (20 fath) depth interval, between 183-327 m (100-179 fath). Achievement of this sampling goal was constrained by the steep bathymetry of many areas, and time.

Methods for determining species compositions follow those of Nagtegaal et al. (1986). Estimates of catch weight were provided by the vessel master. The catch was subsampled for species composition from the conveyor, and the proportions derived were applied to the estimate of weight of total catch for the haul. Catch weights were recorded in pounds and later converted to kilograms. Catch estimates were checked against known landing weights.

Biological samples for major species consisted of 100 specimens for which length/sex/maturity/double otoliths (L/S/M/O) were obtained. Another 100 L/S/M were obtained for each additional sample in a depth interval. Fork lengths were measured to the nearest cm. Forty-eight pairs of Sebastes alutus gills from shallow and deep tows at each locality were also obtained. These gills were preserved for subsequent examination of the copepod parasite Neobrachiella robusta, which has been used for stock identification in S. alutus (Leaman and Kabata 1987). A random sample of 20 S. alutus from the west coast of the Queen Charlotte Islands was also obtained and frozen for taxonomic analysis.

Samples of S. flavidus from Goose Island Gully and off Nootka Sound were frozen for subsequent whole-body parasite screening (Lee et al., in prep.). These samples will be used to determine if a parasite, suitable for use as a natural tag in stock identification, can be found.

RESULTS AND DISCUSSION

Catch

Thirty-eight bottom tows and two midwater tows were completed during the charter (Table 1, Figs. 1-3). S. alutus accounted for 61.2% of the total catch (Table 2). S. reedi, S. entomelas and S. flavidus were also caught in large quantities. The latter two were caught mainly in the two midwater tows; haul 17 off Nootka, and haul 40 in Goose Island Gully (Appendix Table 2, Figs. 1 and 2). Arrowtooth flounder (Atheresthes stomias) and sablefish (Anoplopoma fimbria) were the only non-rockfish species caught in quantity.

Rockfish species composition

S. alutus was consistently the major species of the rockfish catch by bottom trawl, between 183 and 327 m (Table 3, Figs. 4-11). The three cases in which it was not the primary species were: a tow in deep water (328-364 m) off Langara Spit (Fig. 4), where S. aleutianus was the major species caught; a tow in the 219-254 m interval off Langara Spit, where S. zacentrus predominated; and a tow in the 219-254 m interval off Rennell Sound (Fig. 5), where S. reedi and S. proriger had higher catch rates.

S. brevispinis, S. entomelas, S. flavidus and S. proriger usually exhibited highest catch rates in the shallowest depth stratum fished, and a general trend of decreased catch rate with depth in all areas. S. reedi, S. proriger and S. zacentrus generally showed decreased catch rates with depth off the west coasts of the Queen Charlotte and Vancouver islands (Figs. 4-6, 10 and 11). S. diploproa and S. crameri catch rates increased with depth off the west coast of Vancouver Island (Figs. 10 and 11) and in Moresby and Mitchell's gullies (Figs. 7 and 8), but were not caught in quantity in the Goose Island Gully (Fig. 9) or off the west coast of the Queen Charlotte Islands (Figs 4-6). S. aleutianus, S. borealis, and Sebastolobus alascanus all exhibited a general trend of increased catch rate with depth, the former two usually showing maximum catch rates at the deepest intervals sampled. Depths below 273 m were not fished off the west coast of Vancouver Island, so the importance of S. aleutianus and S. borealis could not be assessed. A previous survey of the area encountered small amounts of these species, but only in quantity at depths below 330 m (Leaman et al. 1988).

The two midwater tow catches were primarily S. entomelas (72% of the catch off Nootka, 54% in Goose Island Gully) and S. flavidus. The Goose Island Gully catch also contained small proportions of S. paucispinis and S. pinniger.

Biological Samples

A total of 7,857 rockfish were sampled for length and sex, and 3,728 of these were assigned gonad maturity stages and had otoliths collected (Table 4). Seventeen samples of 48 pairs of gills were taken from Pacific ocean perch to document the incidence and prevalence of the copepod parasite Neobrachiella robusta (Table 5). Twenty whole S. alutus were frozen to measure morphometric differences between two body forms observed off the west coast of the Queen Charlotte Islands. Two samples of twenty S. flavidus were taken for whole-body parasite assay. The results from these collections will be presented elsewhere.

Length frequency data from these samples are generally unremarkable. Due to the determinate growth pattern of rockfishes many age-classes may be represented in each length class, particularly at ages above 25 years (Archibald et al. 1981). There were no instances of large proportions of very small fish in any of the samples, and thus no evidence of recruitment of strong year-classes in the 1983-1986 period.

Sex ratio for S. alutus was nearly equal for most of the stocks (Table 6).

Proportions of the samples at different reproductive development stages were fairly constant throughout the areas, with the exception that there were greater proportions of immature fish recorded from Moresby and Goose Island gullies, and southwest Vancouver Island (Table 7).

Off the west coast of the Queen Charlotte Islands the majority of female S. alutus sampled were in the small maturing stage, and most of the males were also maturing. All three areas (Langara, Rennell Sound and Anthony Island) had very few (<3%) immature fish in the samples.

Proportions of immature S. alutus in Moresby Gully were 10.2% for females and 9.0% for males. Proportions of immatures in Mitchell's Gully were 5.4% for females and 6.7% for males. In Goose Island Gully, proportions of immatures were extremely high at 44.2% for females and 37.0% for males. The majority of the adult females were in the maturing and resting stages (stages 2, 3 and 7) and the adult males were predominantly maturing (stage 2). The higher proportions of immature fish are correlated with a greater number of small fish (<35 cm) in the length frequency samples from these areas.

Off the west coast of Vancouver Island, the bulk of the fish sampled for both sexes were in the maturing stages. Percentage of immature fish sampled was highest off southwest Vancouver Island (8.9% for females and 17.5% for males) and lowest off Nootka (1.9% for females and 4.3% for males).

L/S/M/O and L/S/M samples were taken for S. entomelas (Table 8), S. flavidus (Table 9), S. aleutianus (Table 10), S. reedi (Table 11) and S. brevispinis (Table 12) when these species were caught in sufficient quantities. A L/S/O sample of Pacific hake, Merluccius productus, was taken from haul 40 (Table 13).

ACKNOWLEDGEMENTS

We would like to thank Capt. Dave Clattenburg and the crew of the F/V OCEAN SELECTOR for their cooperation and assistance in the completion of the cruise. Thanks to Rick Stanley and Deanna Lee, whose efforts were instrumental in the successful completion of the cruise. Ray Foucher reviewed a draft of the report and provided thoughtful criticisms.

LITERATURE CITED

- Anon. 1989. Pacific groundfish management plan. Manuscript prepared by Offshore Division, Fisheries Branch, Department of Fisheries and Oceans. Vancouver, B.C. 27 p.
- Archibald, C. P., W. Shaw and B. M. Leaman. 1981. Growth and mortality estimates of rockfishes (Scorpaenidae) from B.C. coastal waters, 1977-1979. Can. Tech. Rep. Fish. Aquat. Sci. 1048: 57 p.
- Leaman, B. M. 1985. Slope rockfishes. In: A.V. Tyler and G.A. McFarlane [eds.]. Groundfish stock assessments for the west coast of Canada in 1984 and recommended yield options for 1985. Can MS Rep. Fish. Aquat. Sci. 1813: 250-307.
- Leaman, B. M., G. E. Gillespie, D. A. Nagtegaal and R. D. Stanley. 1988. Biomass survey of rockfish stocks off the southwest coast of Vancouver Island, September 8-22, 1985 (M/V HOWE BAY). Can. Tech. Rep. Fish. Aquat. Sci. 1611: 99 p.
- Leaman, B. M. and Z. Kabata. 1987. Neobrachiella robusta (Wilson 1912) (Copepoda: Lernaepodidae) as a tag for identification of stocks of its host, Sebastes alutus (Gilbert 1890) (Pisces: Teleostei). Can. J. Zool. 65(11): 2579-2582.

- Lee, D., D. Whitaker and R. D. Stanley. (In prep.) A preliminary examination of the parasite fauna of yellowtail rockfish, Sebastes flavidus. Can. MS Rep. Fish. Aquat. Sci.
- Nagtegaal, D. A., R. D. Stanley and B. M. Leaman. 1986. Catches and trawl locations of the R/V G.B. REED and F/V EASTWARD HO during Pacific ocean perch assessment cruise to Queen Charlotte Sound, August-September, 1984. Can. Data Rep. Fish. Aquat. Sci. 611: 109 p.

Table 1. Haul numbers and total effort by area and gear, F/V OCEAN SELECTOR, July 12 to 30, 1989. All hauls are bottom trawl, unless designated as midwater trawl (MW).

Locality	Haul #	Effort (hr)
Moresby Gully	1-8, 34-36	16.46
Mitchell's Gully	9-11	7.68
Quatsino	12-15	7.72
Nootka	16	1.00
Nootka (MW)	17	1.67
Southwest Vancouver Island	18-22	10.50
Langara	23-27	9.75
Rennell Sound	28-31	10.95
Anthony Island	32-33	2.77
Goose Island Gully	37-39	6.90
Goose Island Gully (MW)	40	3.17
Totals	40	78.57

Table 2. Catch weights and percent of total catch by species, F/V OCEAN SELECTOR, July 12 to 30, 1989. Other species includes all those which represented less than 1% of the total catch. See Table 14 for common names of species.

Species	Weight (kg)	Percent of total
<u>Sebastes alutus</u>	165,621	61.2
<u>S. reedi</u>	16,404	6.1
<u>S. entomelas</u>	16,296	6.0
<u>S. flavidus</u>	11,492	4.2
<u>S. aleutianus</u>	8,414	3.1
<u>S. zacentrus</u>	7,345	2.7
<u>S. diploproa</u>	6,344	2.3
<u>S. brevispinis</u>	5,592	2.1
<u>Atheresthes stomias</u>	5,118	1.9
<u>S. proriger</u>	4,639	1.7
<u>Anoplopoma fimbria</u>	4,617	1.7
<u>Sebastolobus alascanus</u>	4,209	1.6
Other species	14,608	5.4
Total catch	270,966	100.0

Table 3. Mean rockfish catch rates (kg/hr) by area and depth stratum, F/V OCEAN SELECTOR, July 12 to 30, 1989. Means are calculated as averages of catch rates between tows within an area and depth stratum. Midwater tows are presented separately (MW).

Area	Langara			
	219-254	255-273	274-327	328-364
Depth (m)				
<u>S. aleutianus</u>	0	11.38	130.96	7115.05
<u>S. alutus</u>	554.78	2436.84	1827.87	1746.05
<u>S. babcocki</u>	0	34.25	14.78	0
<u>S. borealis</u>	0	0	276.05	0
<u>S. brevispinis</u>	110.23	29.68	0	0
<u>S. crameri</u>	0	0	14.78	0
<u>S. diploproa</u>	0	0	0	0
<u>S. elongatus</u>	0	0	0	0
<u>S. entomelas</u>	0	3.88	0	0
<u>S. flavidus</u>	0	0	0	0
<u>S. helvomaculatus</u>	0	0	14.78	0
<u>S. paucispinis</u>	0	0	0	0
<u>S. pinniger</u>	0	0	0	0
<u>S. proriger</u>	0	3.88	0	0
<u>S. reedi</u>	4.41	11.38	0	0
<u>S. variegatus</u>	79.14	0	0	0
<u>S. zacentrus</u>	652.96	253.49	210.60	0
<u>Sebastolobus alascanus</u>	21.87	32.68	109.14	79.05
No. of Tows	1	2	1	1

Table 3 (cont'd)

Area	Rennell Sound			Anthony Island	
	219-254	255-273	274-327	255-273	274-327
Depth (m)					
<u>S. aleutianus</u>	0	18.77	85.82	0	95.68
<u>S. alutus</u>	415.65	1589.13	4208.15	4825.76	1553.18
<u>S. babcocki</u>	0	2.92	28.76	66.91	71.93
<u>S. borealis</u>	0	18.77	0	0	23.80
<u>S. brevispinis</u>	185.00	58.52	0	84.05	11.93
<u>S. crameri</u>	0	0	0	0	0
<u>S. diploproa</u>	0	0	0	0	0
<u>S. elongatus</u>	0	0	0	0	0
<u>S. entomelas</u>	138.77	0	0	17.19	0
<u>S. flavidus</u>	0	0	0	0	0
<u>S. helvomiculatus</u>	46.29	27.01	0	17.19	23.80
<u>S. paucispinis</u>	0	14.60	0	0	0
<u>S. pinniger</u>	0	0	0	0	0
<u>S. proriger</u>	1501.15	17.65	0	133.76	18.17
<u>S. reedi</u>	5634.00	140.53	0	133.76	0
<u>S. variegatus</u>	0	0	0	0	0
<u>S. zacentrus</u>	0	0	0	352.34	48.17
<u>Sebastolobus alascanus</u>	23.17	20.53	185.57	100.34	101.93
No. of Tows	1	2	1	1	1

Table 3 (cont'd)

Area Depth (m)	Moresby Gully				
	183-218	219-254	255-273	274-327	328-364
<u>S. aleutianus</u>	0	0	7.30	0	286.38
<u>S. alutus</u>	2796.80	2177.24	1048.78	14681.71	2272.46
<u>S. babcocki</u>	15.05	14.53	53.56	92.02	179.29
<u>S. borealis</u>	0	0	0	123.07	366.89
<u>S. brevispinis</u>	1353.80	714.83	3.61	2.53	0
<u>S. crameri</u>	0	0	0	8.55	18.28
<u>S. diploproa</u>	0	0	76.55	29.22	0
<u>S. elongatus</u>	0	0	0	0	0
<u>S. entomelas</u>	59.30	0	0	0	0
<u>S. flavidus</u>	342.05	0	0	0	0
<u>S. helvomiculatus</u>	0	0	1.30	0	0
<u>S. paucispinis</u>	0	26.30	0	142.25	0
<u>S. pinniger</u>	119.30	0	0	0	0
<u>S. proriger</u>	0	0	0	0	0
<u>S. reedi</u>	0	21.92	1.94	157.82	0
<u>S. variegatus</u>	0	0	0	0	0
<u>S. zacentrus</u>	0	131.06	3.83	204.56	0
<u>Sebastolobus alascanus</u>	0	40.78	67.70	177.58	277.27
No. of Tows	1	2	3	4	1

Table 3 (cont'd)

Area Depth (m)	Mitchell's Gully	
	219-254	255-273
<u>S. aleutianus</u>	0	4.02
<u>S. alutus</u>	1017.32	648.11
<u>S. babcocki</u>	21.41	4.47
<u>S. borealis</u>	0	0
<u>S. brevispinis</u>	17.32	25.98
<u>S. crameri</u>	0	0
<u>S. diploproa</u>	0	53.01
<u>S. elongatus</u>	0	0
<u>S. entomelas</u>	0	0
<u>S. flavidus</u>	8.69	0
<u>S. helvomaculatus</u>	17.32	0
<u>S. paucispinis</u>	0	0
<u>S. pinniger</u>	0	0
<u>S. proriger</u>	0	16.38
<u>S. reedi</u>	21.41	107.43
<u>S. variegatus</u>	0	0
<u>S. zacentrus</u>	0	23.33
<u>Sebastolobus alascanus</u>	43.23	17.50
No. of Tows	1	2

Table 3 (cont'd)

Area Depth (m)	Goose Island Gully			
	110-145(MW)	146-182	219-254	255-273
<u>S. aleutianus</u>	0	0	0	20.05
<u>S. alutus</u>	0	477.76	1316.81	606.72
<u>S. babcocki</u>	0	135.16	120.86	20.05
<u>S. borealis</u>	0	0	0	5.83
<u>S. brevispinis</u>	0	52.26	42.62	0
<u>S. crameri</u>	0	0	0	0
<u>S. diploproa</u>	0	0	0	0
<u>S. elongatus</u>	0	0	0	0
<u>S. entomelas</u>	3468.37	30.28	0	0
<u>S. flavidus</u>	2869.00	0	0	0
<u>S. helvomaculatus</u>	0	0	0	0
<u>S. paucispinis</u>	36.37	60.51	0	11.61
<u>S. pinniger</u>	0	0	0	0
<u>S. proriger</u>	18.05	5.55	0	0
<u>S. reedi</u>	0	0	0	0
<u>S. variegatus</u>	0	0	0	0
<u>S. zacentrus</u>	0	0	0	0
<u>Sebastolobus alascanus</u>	0	33.03	78.29	22.72
No. of Tows	1	1	1	1

Table 3 (cont'd)

Area Depth (m)	Quatsino		Nootka	
	183-218	219-254	146-182(MW)	255-273
<u>S. aleutianus</u>	0	0	0	0
<u>S. alutus</u>	1311.07	2028.31	0	849.50
<u>S. babcocki</u>	4.05	5.63	0	16.05
<u>S. borealis</u>	0	0	0	0
<u>S. brevispinis</u>	103.14	0	0	0
<u>S. crameri</u>	0	112.99	0	8.05
<u>S. diploproa</u>	0	1363.07	0	409.05
<u>S. elongatus</u>	4.05	0	0	0
<u>S. entomelas</u>	0	0	2292.65	0
<u>S. flavidus</u>	0	0	1159.85	0
<u>S. helvomaculatus</u>	4.05	10.52	0	0
<u>S. paucispinis</u>	64.45	0	0	0
<u>S. pinniger</u>	0	0	0	0
<u>S. proriger</u>	240.89	0	0	0
<u>S. reedi</u>	233.32	0	0	0
<u>S. variegatus</u>	0	0	0	0
<u>S. zacentrus</u>	428.36	45.20	0	8.05
<u>Sebastolobus alascanus</u>	20.94	79.89	0	102.05
No. of Tows	2	2	1	1

Table 3 (cont'd)

Area Depth (m)	Southwest Vancouver Island		
	183-218	219-254	255-273
<u>S. aleutianus</u>	0	0	0
<u>S. alutus</u>	750.87	548.18	998.05
<u>S. babcocki</u>	9.38	30.63	10.55
<u>S. borealis</u>	0	0	0
<u>S. brevispinis</u>	6.58	0	0
<u>S. crameri</u>	0	0	92.55
<u>S. diploproa</u>	95.63	136.31	389.55
<u>S. elongatus</u>	86.24	11.28	0
<u>S. entomelas</u>	1.31	0	0
<u>S. flavidus</u>	0	0	0
<u>S. helvomaculatus</u>	14.37	13.99	10.55
<u>S. paucispinis</u>	0	0	0
<u>S. pinniger</u>	0	0	0
<u>S. proriger</u>	54.41	0	0
<u>S. reedi</u>	489.84	0	0
<u>S. variegatus</u>	0	0	0
<u>S. zacentrus</u>	66.66	52.70	5.05
<u>Sebastolobus alascanus</u>	32.53	61.21	41.05
No. of Tows	3	1	1

Table 4. Summary of rockfish biological samples taken on the F/V OCEAN SELECTOR, July 12 to 30, 1989. (Sample types: L/S/M = length, sex and maturity stage; L/S/M/O = length, sex, maturity stage and double otoliths; MORPH = whole fish frozen for morphometric analyses; PARASITE = whole fish frozen for parasite assay; RAKER = gill raker parasite sample.)

Species	Area	Sample type	N
<u>Sebastes</u>	Langara	L/S/M	99
<u>aleutianus</u>	Langara	L/S/M/O	100
Total			199
<u>S. alutus</u>	Langara	L/S/M	431
	Langara	L/S/M/O	400
	Langara	RAKER	96
	Rennell Sound	L/S/M	314
	Rennell Sound	L/S/M/O	300
	Rennell Sound	RAKER	96
	Rennell Sound	MORPH	20
	Anthony Island	L/S/M	199
	Anthony Island	L/S/M/O	200
	Anthony Island	RAKER	96
	Moresby Gully	L/S/M	1001
	Moresby Gully	L/S/M/O	900
	Moresby Gully	RAKER	144
	Mitchell's Gully	L/S/M	198
	Mitchell's Gully	L/S/M/O	200
	Mitchell's Gully	RAKER	96
	Goose Island Gully	L/S/M	421
	Goose Island Gully	L/S/M/O	400
	Goose Island Gully	RAKER	96
	Quatsino	L/S/M	431
	Quatsino	L/S/M/O	400
	Quatsino	RAKER	96
	Nootka	L/S/M	132
	Nootka	L/S/M/O	100
	SWVI	L/S/M	428
	SWVI	L/S/M/O	300
	SWVI	RAKER	96
Total			6,775
<u>S. brevispinis</u>	Moresby Gully	L/S/M	116
	Moresby Gully	L/S/M/O	100
Total			216

Table 4 (cont'd)

Species	Area	Sample type	N
<u>S. entomelas</u>	Goose Island Gully	L/S/M/O	10
	Nootka	L/S/M/O	18
	Total		28
<u>S. flavidus</u>	Goose Island Gully	L/S/M	100
	Goose Island Gully	L/S/M/O	100
	Goose Island Gully	PARASITE	20
	Nootka	L/S/M	105
	Nootka	L/S/M/O	100
	Nootka	PARASITE	20
	Total		445
<u>S. reedi</u>	Rennell Sound	L/S/M	94
	Rennell Sound	L/S/M/O	100
	Total		194
Grand Total All Rockfish			7,857

Table 5. Summary of sample data for gill raker collections from Pacific ocean perch, S. alutus, F/V OCEAN SELECTOR, July 12 to 30, 1989. Depths presented are the modal depth for the tow (Appendix Table 2).

Haul	Area	Depth (m)	N
1	Moresby Gully	275	48
3	Moresby Gully	320	48
4	Moresby Gully	338	48
9	Mitchell's Gully	245	48
10	Mitchell's Gully	260	48
12	Quatsino	210	48
13	Quatsino	247	48
18	Southwest Vancouver Island	283	48
19	Southwest Vancouver Island	229	48
24	Langara	262	48
25	Langara	238	48
28	Rennell Sound	223	48
30	Rennell Sound	309	48
32	Anthony Island	256	48
33	Anthony Island	320	48
37	Goose Island Gully	165	48
39	Goose Island Gully	274	48
Total			816

Table 6. Summary of length frequency and maturity of Sebastes alutus collected aboard the F/V OCEAN SELECTOR, July 12 to 30, 1989. See Appendix Table 3 for maturity code descriptions.

Stock	260789		Langara		260789	
Date (DDMMYY)	260789		260789		260789	
Depth (m)	262		238		256	
Set	24		25		26	
	M	F	M	F	M	F
Length (cm)						
17	-	-	-	-	-	-
18	-	-	-	-	-	-
19	-	-	-	-	-	-
20	-	-	-	-	-	-
21	-	-	-	-	-	-
22	-	-	-	-	-	-
23	-	-	-	-	-	-
24	-	-	-	-	-	-
25	-	-	-	-	-	-
26	1	-	1	-	-	-
27	0	-	0	-	-	-
28	0	-	0	-	-	-
29	0	-	0	-	1	-
30	0	-	1	-	0	-
31	0	-	0	1	1	-
32	0	1	1	1	2	2
33	2	2	4	1	4	1
34	12	3	9	3	5	8
35	12	12	15	10	23	11
36	17	14	11	14	25	19
37	20	16	22	13	16	14
38	19	11	20	16	16	11
39	8	14	9	17	10	18
40	6	9	6	8	4	11
41	3	16	0	6	-	6
42	1	2	1	1	-	1
43	1	4	-	2	-	0
44	-	2	-	4	-	2
45	-	1	-	2	-	1
46	-	1	-	1	-	-
47	-	-	-	-	-	-
48	-	-	-	-	-	-
49	-	-	-	-	-	-
50	-	-	-	-	-	-
51	-	-	-	-	-	-
Total	102	108	100	100	107	105
Maturities						
1	1	0	2	0	1	0
2	49	46	46	50	44	55
3	0	0	0	0	0	0
4	0	0	0	0	0	0
5	0	0	0	0	0	0
6	0	0	0	0	0	0
7	1	3	0	2	0	0
9	0	0	0	0	0	0

Table 6 (cont'd)

Stock Date (DDMMYY) Depth (m) Set	Langara 260789 311 27		Total for Langara	
	M	F	M	F
Length (cm)				
16	-	-	-	-
17	-	-	-	-
18	-	-	-	-
19	-	-	-	-
20	-	-	-	-
21	-	-	-	-
22	-	-	-	-
23	-	-	-	-
24	-	-	-	-
25	-	-	-	-
26	1	-	3	-
27	0	-	0	-
28	1	-	1	-
29	0	1	1	1
30	1	0	2	0
31	2	2	3	3
32	3	1	6	5
33	6	3	16	7
34	10	4	36	18
35	19	12	69	45
36	7	10	60	57
37	17	17	75	60
38	9	15	64	53
39	9	23	36	72
40	2	12	18	40
41	2	10	5	38
42	2	5	4	9
43	-	2	1	8
44	-	0	-	8
45	-	1	-	5
46	-	-	-	2
47	-	-	-	-
48	-	-	-	-
49	-	-	-	-
50	-	-	-	-
51	-	-	-	-
Total	91	118	400	431
Maturities				
1	1	0	5	0
2	41	58	180	209
3	0	0	0	0
4	0	0	0	0
5	0	0	0	0
6	0	0	0	0
7	0	0	1	5
9	0	0	0	0

Table 6 (cont'd)

Stock Date (DDMMYY)	270789		Rennell Sound		270789		Total for Rennell Sound	
	223		278		309			
Depth (m)	28		29		30			
Set	M	F	M	F	M	F	M	F
Length (cm)								
16	-	-	-	-	-	-	-	-
17	-	-	-	-	-	-	-	-
18	-	-	-	-	-	-	-	-
19	-	-	-	-	-	-	-	-
20	-	-	-	-	-	-	-	-
21	-	-	-	-	-	-	-	-
22	-	-	-	-	-	-	-	-
23	-	-	-	-	-	-	-	-
24	-	-	-	-	-	-	-	-
25	-	-	-	-	-	-	-	-
26	-	-	-	-	-	-	-	-
27	-	-	-	-	-	-	-	-
28	-	-	-	-	-	-	-	-
29	-	-	-	-	-	-	-	-
30	-	-	-	1	-	1	-	2
31	-	-	-	0	-	0	-	0
32	-	1	1	0	-	0	1	1
33	-	0	0	0	-	0	0	0
34	-	1	0	2	-	0	0	3
35	4	0	1	0	1	3	6	3
36	4	2	3	1	6	1	13	4
37	6	6	10	3	11	4	27	13
38	8	8	9	6	6	7	23	21
39	16	5	31	12	13	10	60	27
40	20	12	9	9	6	15	35	36
41	12	25	7	18	7	15	26	58
42	4	16	3	19	5	14	12	49
43	3	9	0	11	1	25	4	45
44	1	8	0	8	-	13	1	29
45	0	6	0	8	-	14	0	28
46	1	9	0	13	-	7	1	29
47	-	11	0	9	-	10	0	30
48	-	4	0	5	-	4	0	13
49	-	4	1	1	-	1	1	6
50	-	0	-	-	-	-	-	0
51	-	1	-	-	-	-	-	1
Total	79	128	75	126	56	144	210	398
Maturities								
1	0	0	0	1	0	1	0	2
2	39	56	37	57	26	73	102	186
3	0	0	0	0	0	0	0	0
4	0	0	0	0	0	0	0	0
5	0	0	0	0	0	0	0	0
6	0	0	0	0	0	0	0	0
7	0	5	0	5	0	0	0	10
9	0	0	0	0	0	0	0	0

Table 6 (cont'd)

Stock Date (DDMMYY) Depth (m) Set	Anthony Island				Total for Anthony Island	
	280789		280789			
	142 32	F	220 33	F	M	F
Length (cm)						
16	-	-	-	-	-	-
17	-	-	-	-	-	-
18	-	-	-	-	-	-
19	-	-	-	-	-	-
20	-	-	-	-	-	-
21	-	-	-	-	-	-
22	-	-	-	-	-	-
23	-	-	-	-	-	-
24	-	-	-	-	-	-
25	-	-	-	-	-	-
26	-	-	-	-	-	-
27	-	-	-	-	-	-
28	-	-	-	-	-	-
29	-	-	-	-	-	-
30	-	-	-	1	-	1
31	-	-	-	1	-	1
32	1	-	-	0	1	0
33	0	1	1	0	1	1
34	0	0	0	1	0	1
35	2	2	5	3	7	5
36	2	4	10	2	12	6
37	5	5	10	8	15	13
38	11	5	6	8	17	13
39	10	9	8	8	18	17
40	10	13	18	16	28	29
41	6	18	13	5	19	23
42	8	19	16	15	24	34
43	8	16	5	10	13	26
44	4	16	3	13	7	29
45	-	13	-	5	-	18
46	-	6	-	6	-	12
47	-	1	-	1	-	2
48	-	4	-	0	-	4
49	-	1	-	1	-	2
50	-	-	-	-	-	-
51	-	-	-	-	-	-
Total	67	133	95	104	162	237
Maturities						
1	2	0	0	0	2	0
2	27	70	54	41	81	111
3	0	0	0	0	0	0
4	0	0	0	0	0	0
5	0	0	0	0	0	0
6	0	0	0	0	0	0
7	0	1	0	5	0	6
9	0	0	0	0	0	0

Table 6 (cont'd)

Stock Date (DDMMYY)	Moresby Gully							
	140789		140789		140789		140789	
Depth (m)	275		302		320		338	
Set	1		2		3		4	
	M	F	M	F	M	F	M	F
Length (cm)								
17	-	-	-	-	-	-	-	-
18	-	-	-	-	-	-	-	-
19	-	-	-	-	-	-	-	-
20	-	-	-	-	-	-	-	-
21	-	-	-	-	-	-	-	-
22	-	-	-	-	-	-	-	-
23	-	-	-	-	-	-	-	-
24	-	-	-	-	-	-	-	-
25	-	-	-	-	-	-	-	-
26	-	1	-	-	-	-	-	-
27	-	0	-	-	-	-	-	-
28	-	0	-	-	-	-	-	-
29	-	0	-	-	-	-	-	-
30	-	0	-	-	-	-	-	-
31	-	0	-	-	-	-	-	-
32	-	0	-	1	-	-	1	-
33	1	0	1	1	-	1	0	-
34	1	1	3	1	4	1	0	-
35	2	7	12	2	5	4	1	-
36	14	1	7	7	11	5	3	-
37	16	4	22	20	9	8	7	2
38	24	12	9	13	14	14	7	1
39	20	8	12	18	16	20	26	7
40	9	19	10	23	18	21	43	4
41	15	11	4	16	7	19	47	12
42	8	4	1	1	0	12	35	2
43	0	3	2	0	4	7	10	5
44	1	6	-	1	1	3	2	2
45	2	5	-	5	-	6	1	0
46	-	4	-	4	-	8	-	5
47	-	1	-	4	-	7	-	2
48	-	1	-	2	-	0	-	-
49	-	-	-	-	-	1	-	-
50	-	-	-	-	-	-	-	-
51	-	-	-	-	-	-	-	-
Total	113	88	83	119	89	137	183	42
Maturities								
1	2	3	1	3	1	0	0	0
2	79	1	38	0	36	0	78	0
3	0	42	0	52	0	49	0	18
4	0	0	0	0	0	0	0	0
5	0	0	0	0	0	0	0	0
6	0	0	0	0	0	0	0	0
7	0	15	0	6	0	14	0	4
9	0	0	0	0	0	0	0	0

Table 6 (cont'd)

Stock		Moresby Gully					
Date (DDMMYY)		140789		150789		150789	
Depth (m)		265		294		274	
Set		5		6		7	
		M	F	M	F	M	F
Length (cm)							
16		-	-	-	-	-	-
17		-	-	-	-	-	-
18		-	-	-	-	-	-
19		-	-	-	-	-	-
20		-	-	1	-	-	-
21		-	-	0	-	-	-
22		2	-	0	-	-	-
23		1	-	0	-	-	-
24		0	1	0	-	-	-
25		0	0	0	-	2	-
26		1	0	0	-	6	2
27		0	0	0	-	2	2
28		0	1	0	-	3	0
29		0	0	0	-	5	4
30		1	2	0	1	8	3
31		1	7	2	1	4	5
32		4	4	2	0	5	4
33		5	2	4	0	5	2
34		3	1	1	2	7	9
35		7	4	2	3	6	6
36		14	10	11	3	10	7
37		18	5	13	4	7	8
38		17	11	15	13	10	6
39		8	16	4	9	10	7
40		5	20	14	4	10	7
41		8	10	23	8	6	5
42		7	10	21	3	5	4
43		6	4	13	9	3	1
44		0	4	1	5	4	3
45		0	2	-	5	1	1
46		1	2	-	3	0	0
47		-	2	-	3	0	0
48		-	0	-	-	0	1
49		-	1	-	-	1	-
50		-	-	-	-	-	-
51		-	-	-	-	-	-
Total		109	119	127	76	120	87
Maturities							
1		5	7	2	1	19	7
2		35	0	52	12	44	0
3		0	43	14	0	0	20
4		0	0	0	0	1	0
5		0	0	0	0	0	0
6		0	0	0	0	0	0
7		0	10	0	19	1	8
9		0	0	0	0	0	0

Table 6 (cont'd)

Stock Date (DDMMYY) Depth (m) Set	Moresby Gully				Total for Moresby Gully	
	280789		280789			
	219 34		282 36			
	M	F	M	F	M	F
Length (cm)						
16	-	-	-	-	-	-
17	-	-	-	-	-	-
18	-	-	-	-	-	-
19	-	-	-	-	-	-
20	-	-	-	-	1	-
21	-	-	-	-	0	-
22	1	1	-	-	3	1
23	0	1	-	-	1	1
24	2	2	-	-	2	3
25	0	4	-	-	2	4
26	3	1	-	-	10	4
27	1	0	1	1	4	3
28	4	0	1	0	8	1
29	3	1	0	2	8	7
30	2	4	0	3	11	13
31	2	4	7	2	16	19
32	3	7	2	5	17	21
33	9	8	4	8	29	22
34	6	6	4	4	29	25
35	0	6	5	3	40	35
36	6	5	5	3	81	41
37	10	6	7	12	109	69
38	11	4	15	12	122	86
39	9	8	11	3	116	96
40	6	12	12	11	127	121
41	2	13	9	8	121	102
42	2	14	8	6	87	56
43	0	7	7	2	45	38
44	1	3	6	1	16	28
45	-	4	2	2	6	30
46	-	1	4	1	5	28
47	-	1	1	1	1	21
48	-	-	-	1	0	5
49	-	-	-	-	1	2
50	-	-	-	-	-	-
51	-	-	-	-	-	-
Total	83	123	111	91	1018	882
Maturities						
1	8	8	6	17	44	46
2	31	47	38	30	431	90
3	0	0	0	0	14	224
4	0	0	0	0	1	0
5	0	0	0	0	0	0
6	0	0	0	0	0	0
7	0	6	0	9	1	91
9	0	0	0	0	0	0

Table 6 (cont'd)

Stock Date (DDMMYY)	160789		Mitchell's Gully				Total for Mitchell's Gully	
	245		160789		160789			
Depth (m)	9		10		11			
Set	M	F	M	F	M	F	M	F
Length (cm)								
16	-	-	-	-	-	-	-	-
17	-	-	-	-	-	-	-	-
18	-	-	-	-	-	-	-	-
19	-	-	-	-	-	-	-	-
20	-	-	-	-	-	-	-	-
21	-	-	-	-	-	-	-	-
22	-	-	-	-	1	1	1	1
23	-	-	-	-	1	1	1	1
24	-	-	-	-	1	1	1	1
25	-	-	-	-	0	1	0	1
26	-	-	-	-	1	4	1	4
27	-	1	-	-	4	2	4	3
28	1	0	-	-	1	3	2	3
29	0	0	-	-	1	0	1	0
30	1	0	-	-	0	2	1	2
31	1	2	2	1	3	1	6	4
32	2	1	0	0	2	4	4	5
33	2	0	1	0	8	3	11	3
34	3	3	5	1	7	4	15	8
35	4	3	3	5	7	2	14	10
36	9	8	7	5	8	6	24	19
37	14	3	8	5	11	6	33	14
38	15	11	17	4	14	3	46	18
39	11	16	16	10	12	7	39	33
40	7	8	6	15	8	13	21	36
41	4	10	9	14	4	7	17	31
42	5	8	11	8	1	11	17	27
43	2	7	2	8	3	2	7	17
44	3	9	2	9	-	8	5	26
45	1	9	-	10	-	5	1	24
46	1	10	-	5	-	3	1	18
47	-	5	-	7	-	2	-	14
48	-	1	-	1	-	1	-	3
49	-	-	-	-	-	-	-	-
50	-	-	-	-	-	-	-	-
51	-	-	-	-	-	-	-	-
Total	86	115	89	108	98	103	273	326
Maturities								
1	3	1	1	1	5	7	9	9
2	43	0	42	30	39	47	124	77
3	0	39	0	16	0	1	0	56
4	0	0	0	0	0	0	0	0
5	0	0	0	0	0	0	0	0
6	0	0	0	0	0	0	0	0
7	0	14	0	10	1	0	1	24
9	0	0	0	0	0	0	0	0

Table 6 (cont'd)

Stock Date (DDMMYY)	Goose Island Gully						Total for Goose Island Gully	
Depth (m)	290789		290789		290789			
Set	165 37		219 38		274 39			
	M	F	M	F	M	F	M	F
Length (cm)								
16	-	1	-	-	-	-	-	1
17	1	2	-	-	-	-	1	2
18	3	0	-	-	-	-	3	0
19	0	0	-	-	-	-	0	0
20	0	1	-	-	-	-	0	1
21	3	1	1	1	-	-	4	2
22	1	0	0	1	-	-	1	1
23	7	1	1	5	1	-	9	6
24	6	4	2	7	0	-	8	11
25	6	2	1	3	0	-	7	5
26	0	1	5	2	0	-	5	3
27	4	3	1	3	3	1	8	7
28	2	1	2	9	1	1	5	11
29	1	1	5	6	2	3	8	10
30	2	2	8	4	1	3	11	9
31	7	3	2	8	6	4	15	15
32	7	8	6	9	2	10	15	27
33	6	6	7	7	4	7	17	20
34	10	3	10	2	6	6	26	11
35	2	3	3	1	1	5	6	9
36	2	6	2	3	11	1	15	10
37	11	5	3	4	17	3	31	12
38	8	2	12	5	17	9	37	16
39	7	5	7	5	10	7	24	17
40	7	5	1	7	11	4	19	16
41	8	9	-	5	6	6	14	20
42	3	6	-	6	-	10	3	22
43	1	3	-	8	-	8	1	19
44	1	7	-	6	-	6	1	19
45	-	5	-	4	-	7	-	16
46	-	2	-	3	-	1	-	6
47	-	0	-	1	-	-	-	1
48	-	1	-	-	-	-	-	1
49	-	-	-	-	-	-	-	-
50	-	-	-	-	-	-	-	-
51	-	-	-	-	-	-	-	-
Total	116	99	79	125	99	102	294	326
Maturities								
1	19	18	19	33	12	22	50	73
2	28	28	16	24	41	16	85	68
3	0	0	0	0	0	0	0	0
4	0	0	0	0	0	0	0	0
5	0	0	0	0	0	0	0	0
6	0	0	0	0	0	0	0	0
7	0	7	0	8	0	9	0	24
9	0	0	0	0	0	0	0	0

Table 6 (cont'd)

Stock Date (DDMMYY) Depth (m) Set	170789 210 12		Quatsino 170789 247 13		180789 210 14	
	M	F	M	F	M	F
Length (cm)						
16	-	-	-	-	-	-
17	1	-	-	-	-	-
18	0	-	-	-	1	1
19	0	-	-	-	0	0
20	0	-	-	-	1	0
21	0	-	-	-	1	0
22	0	-	-	-	1	0
23	0	-	-	-	2	2
24	0	1	-	-	0	1
25	0	1	-	-	2	0
26	0	0	-	-	0	0
27	1	0	-	-	0	1
28	0	0	-	-	0	0
29	0	0	1	-	0	0
30	0	0	0	1	0	0
31	1	0	3	0	0	0
32	0	0	7	4	1	0
33	6	1	4	3	2	1
34	5	0	21	6	3	4
35	8	1	18	14	7	3
36	5	9	12	15	3	5
37	7	8	12	19	11	4
38	10	7	10	5	13	5
39	5	14	9	4	11	5
40	5	13	4	5	8	11
41	4	20	6	3	5	11
42	3	17	2	2	4	10
43	1	7	0	9	0	5
44	-	8	3	4	0	7
45	-	14	-	3	1	16
46	-	8	-	2	-	15
47	-	4	-	-	-	12
48	-	4	-	-	-	4
49	-	-	-	-	-	1
50	-	-	-	-	-	-
51	-	-	-	-	-	-
Total	62	137	112	99	77	124
Maturities						
1	2	3	5	3	5	1
2	26	65	44	44	36	56
3	0	0	0	0	0	0
4	0	0	0	0	0	0
5	0	0	0	0	0	0
6	0	0	0	0	0	0
7	0	4	0	4	0	2
9	0	0	0	0	0	0

Table 6 (cont'd)

Stock Date (DDMMYY) Depth (m) Set	Quatsino 180789 256 15		Total for Quatsino	
	M	F	M	F
Length (cm)				
16	-	-	-	-
17	-	-	1	-
18	-	-	1	1
19	-	-	0	0
20	-	-	1	0
21	-	-	1	0
22	-	-	1	0
23	-	-	2	2
24	-	-	0	2
25	-	-	2	1
26	1	-	1	0
27	1	-	2	1
28	0	-	0	0
29	0	-	1	0
30	0	1	0	2
31	0	0	4	0
32	5	1	13	5
33	4	2	16	7
34	5	4	34	14
35	7	4	40	22
36	6	3	26	32
37	11	9	41	40
38	13	2	46	19
39	23	4	48	27
40	24	2	41	31
41	30	6	45	40
42	18	7	27	36
43	8	4	9	25
44	1	5	4	24
45	1	2	2	35
46	-	5	-	30
47	-	-	-	16
48	-	-	-	8
49	-	-	-	1
50	-	-	-	-
51	-	-	-	-
Total	158	61	409	421
Maturities				
1	6	2	18	9
2	63	26	169	191
3	0	0	0	0
4	0	0	0	0
5	0	0	0	0
6	0	0	0	0
7	0	3	0	13
9	0	0	0	0

Table 6 (cont'd)

Stock Date (DDMMYY) Depth (m) Set	Nootka 180789 289 16		Total for Nootka	
	M	F	M	F
Length (cm)				
16	-	-	-	-
17	-	-	-	-
18	-	-	-	-
19	-	-	-	-
20	-	-	-	-
21	-	-	-	-
22	-	-	-	-
23	-	-	-	-
24	-	-	-	-
25	-	-	-	-
26	-	-	-	-
27	-	-	-	-
28	-	-	-	-
29	-	-	-	-
30	1	-	1	-
31	0	3	0	3
32	6	4	6	4
33	9	6	9	6
34	10	5	10	5
35	25	20	25	20
36	13	18	13	18
37	8	21	8	21
38	9	13	9	13
39	7	8	7	8
40	7	12	7	12
41	3	8	3	8
42	3	3	3	3
43	-	4	-	4
44	-	2	-	2
45	-	0	-	0
46	-	2	-	2
47	-	1	-	1
48	-	0	-	0
49	-	1	-	1
50	-	-	-	-
51	-	-	-	-
Total	101	131	101	131
Maturities				
1	2	1	2	1
2	44	53	44	53
3	0	0	0	0
4	0	0	0	0
5	0	0	0	0
6	0	0	0	0
7	0	0	0	0
9	0	0	0	0

Table 6 (cont'd)

Stock Date (DDMMYY) Depth (m) Set	Southwest Vancouver Island					
	190789		190789		190789	
	283		229		183	
	18		19		20	
	M	F	M	F	M	F
Length (cm)						
16	-	-	-	-	-	-
17	-	-	1	-	-	-
18	-	-	0	-	-	1
19	-	-	0	-	3	0
20	-	-	0	-	0	1
21	-	-	0	1	0	0
22	-	-	2	0	0	0
23	-	-	2	1	1	0
24	-	-	2	0	2	2
25	-	-	2	4	7	6
26	-	-	2	2	10	8
27	-	-	4	0	17	5
28	1	-	3	4	10	3
29	0	-	2	0	5	4
30	1	2	5	3	4	2
31	4	2	8	6	3	1
32	8	6	6	10	1	0
33	8	9	13	14	3	2
34	16	9	16	11	0	0
35	22	16	6	6	0	0
36	10	23	12	5	2	0
37	7	13	5	8	1	0
38	4	13	5	4	1	1
39	4	9	2	7	-	0
40	7	5	2	6	-	0
41	0	5	2	8	-	0
42	0	2	0	0	-	0
43	0	3	1	0	-	0
44	0	1	-	4	-	1
45	0	1	-	2	-	-
46	0	1	-	1	-	-
47	0	-	-	-	-	-
48	0	-	-	-	-	-
49	1	-	-	-	-	-
50	-	-	-	-	-	-
51	-	-	-	-	-	-
Total	93	120	103	107	70	37
Maturities						
1	0	0	15	10	0	0
2	42	58	26	48	0	0
3	0	0	0	0	0	0
4	0	0	0	0	0	0
5	0	0	0	0	0	0
6	0	0	0	0	0	0
7	0	0	0	1	0	0
9	0	0	0	0	0	0

Table 6 (cont'd)

Stock Date (DDMMYY) Depth (m) Set	SW Van. Is. 190789 192 21		Total for Southwest Van. Is.		Grand Total	
	M	F	M	F	M	F
Length (cm)						
16	-	-	-	-	-	1
17	-	-	1	-	3	2
18	-	-	0	1	4	2
19	1	-	4	0	4	0
20	0	-	0	1	2	2
21	0	-	0	1	5	3
22	0	-	2	0	8	3
23	0	-	3	1	16	11
24	0	-	4	2	15	19
25	0	1	9	11	20	22
26	0	1	12	11	32	22
27	0	0	21	5	39	19
28	1	0	15	7	31	22
29	1	0	8	4	27	22
30	0	0	10	7	36	36
31	2	1	17	10	61	55
32	5	3	20	19	83	87
33	8	9	32	34	131	100
34	5	7	37	27	187	112
35	7	9	35	31	242	180
36	8	6	32	34	276	221
37	11	13	24	34	363	276
38	7	12	17	30	381	269
39	7	10	13	26	361	323
40	4	10	13	21	309	342
41	4	8	6	21	256	341
42	3	4	3	6	180	242
43	1	7	2	10	82	192
44	1	11	1	17	35	182
45	-	6	0	9	9	165
46	-	4	0	6	7	133
47	-	2	0	2	1	87
48	-	-	0	-	0	34
49	-	-	1	-	3	12
50	-	-	-	-	-	0
51	-	-	-	-	-	1
Total	76	124	342	388	3209	3540
Maturities						
1	6	6	21	16	151	156
2	31	49	99	155	1315	1140
3	0	0	0	0	14	280
4	0	0	0	0	1	0
5	0	0	0	0	0	0
6	0	0	0	0	0	0
7	0	8	0	9	3	182
9	0	0	0	0	0	0

Table 7. Summary of gonad maturity stages of Pacific ocean perch, *S. alutus*, sampled aboard the F/V OCEAN SELECTOR, July 12 to 30, 1989. See Appendix Table 3 for description of maturity codes. (LAN = Langara, RSD = Rennell Sound, ANT = Anthony Island, MOR = Moresby Gully, MIT = Mitchell's Gully, GIG = Goose Island Gully, QUAT = Quatsino, NOOT = Nootka, SWVI = southwest Vancouver Island)

Females

[illegible]

Males

[illegible]

Table 8. Summary of length frequency and maturity of Sebastes entomelas collected aboard the F/V OCEAN SELECTOR, July 12 to 30, 1989. See Appendix Table 3 for maturity code descriptions.

Area	Goose Is. Gully		Nootka		Total	
Date (DDMMYY)	290789		190789			
Depth (m)	133		165			
Set	40		17			
	M	F	M	F	M	F
Length (cm)						
34	-	-	1	-	1	-
35	-	-	1	1	1	1
36	-	-	0	0	0	0
37	-	-	0	0	0	0
38	-	-	0	0	0	0
39	-	-	0	0	0	0
40	-	-	4	2	4	2
41	-	-	0	1	0	1
42	-	-	0	0	0	0
43	-	-	0	2	0	2
44	-	-	1	0	1	0
45	-	-	1	1	1	1
46	2	-	2	1	4	1
47	1	-	-	-	1	0
48	0	-	-	-	0	0
49	2	-	-	-	2	0
50	2	-	-	-	2	0
51	1	-	-	-	1	0
52	0	-	-	-	0	0
53	1	-	-	-	1	0
54	-	-	-	-	-	0
55	-	-	-	-	-	0
56	-	1	-	-	-	1
Total	9	1	10	8	19	9
Maturities						
1	0	0	2	0	2	0
2	9	0	7	4	16	4
3	0	0	0	0	0	0
4	0	0	1	0	1	0
5	0	0	0	0	0	0
6	0	0	0	0	0	0
7	0	1	0	4	0	5
9	0	0	0	0	0	0

Table 9. Summary of length frequency and maturity of Sebastes flavidus collected aboard the F/V OCEAN SELECTOR, July 12 to 30, 1989. See Appendix Table 3 for maturity code descriptions.

Area	Goose Is. Gully		Nootka		Total	
Date (DDMMYY)	290789		190789			
Depth (m)	133		165			
Set	40		17			
	M	F	M	F	M	F
Length (cm)						
37	-	1	-	-	-	1
38	-	0	2	1	2	1
39	1	0	3	2	4	2
40	0	0	1	1	1	1
41	1	1	7	8	8	9
42	2	0	13	5	15	5
43	7	4	10	3	17	7
44	7	3	24	8	31	11
45	13	8	17	9	30	17
46	15	10	18	12	33	22
47	14	7	8	9	22	16
48	11	16	8	5	19	21
49	5	20	6	11	11	31
50	6	11	-	4	6	15
51	1	14	-	5	1	19
52	-	6	-	4	-	10
53	-	8	-	1	-	9
54	-	3	-	-	-	3
55	-	4	-	-	-	4
56	-	0	-	-	-	0
57	-	1	-	-	-	1
Total	83	117	117	88	200	205
Maturities						
1	0	0	0	0	0	0
2	35	23	39	30	74	53
3	0	0	0	0	0	0
4	0	0	17	0	17	0
5	0	0	0	0	0	0
6	0	0	0	0	0	0
7	5	37	1	13	6	50
9	0	0	0	0	0	0

Table 10. Summary of length frequency and maturity of Sebastes aleutianus collected aboard the F/V OCEAN SELECTOR, July 12 to 30, 1989. See Appendix Table 3 for maturity code descriptions.

Area	Langara
Date (DDMMYY)	260789
Depth (m)	338
Set	23

	M	F
Length (cm)		
38	-	1
39	-	0
40	-	2
41	-	1
42	1	2
43	6	3
44	6	9
45	17	13
46	14	20
47	15	11
48	13	18
49	6	10
50	2	10
51	3	5
52	1	2
53	-	1
54	-	1
55	-	1
56	-	0
57	-	1
58	-	0
59	-	1
Total	84	112

Maturities

1	0	0
2	45	50
3	0	0
4	0	0
5	1	0
6	0	0
7	1	3
9	0	0

Table 11. Summary of length frequency and maturity of Sebastes reedi collected aboard the F/V OCEAN SELECTOR, July 12 to 30, 1989. See Appendix Table 3 for maturity code descriptions.

Area Rennell Sound
Date (DDMMYY) 270789
Depth (m) 223
Set 28

	M	F
Length (cm)		
31	-	1
32	-	1
33	-	0
34	-	1
35	-	1
36	-	0
37	-	0
38	3	2
39	3	5
40	12	10
41	17	18
42	11	16
43	7	16
44	11	19
45	7	19
46	1	10
47	-	2
48	-	1
Total	72	122

Maturities

1	0	1
2	24	7
3	0	0
4	0	0
5	0	0
6	0	0
7	7	61
9	0	0

Table 12. Summary of length frequency and maturity of *Sebastes brevispinis* collected aboard the F/V OCEAN SELECTOR, July 12 to 30, 1989. See Appendix Table 3 for maturity code descriptions.

Area	Moresby
Date (DDMMYY)	280789
Depth (m)	219
Set	34

Length (cm)	M	F
37	-	1
38	-	0
39	-	2
40	-	2
41	-	1
42	1	0
43	0	0
44	0	1
45	2	1
46	1	2
47	5	4
48	15	0
49	14	8
50	21	10
51	12	16
52	13	8
53	14	12
54	6	7
55	5	3
56	2	7
57	3	7
58	0	4
59	1	2
60	-	2
61	-	0
62	-	0
63	-	0
64	-	1

Total	115	101
-------	-----	-----

Maturities

1	0	0
2	31	2
3	0	0
4	0	1
5	0	3
6	0	2
7	28	33
9	0	0

Table 13. Summary of length frequency of Pacific hake, Merluccius productus, collected aboard the F/V OCEAN SELECTOR, July 12 to 30, 1989.

Area Goose Is. Gully
Date (DDMMYY) 290789
Depth (m) 133
Set 40

Length (cm)	M	F
47	1	-
48	0	1
49	0	1
50	0	0
51	4	3
52	1	2
53	1	2
54	1	1
55	0	7
56	0	11
57	0	3
58	1	7
59	0	12
60	0	12
61	1	18
62	-	11
63	-	14
64	-	5
65	-	2
66	-	8
67	-	5
68	-	3
69	-	2
70	-	2
71	-	1
Total	10	133

Table 14. Scientific and common names for species encountered on the F/V OCEAN SELECTOR, July 12 to 30, 1989.

Rockfish

Rougheye rockfish	<u>Sebastes aleutianus</u>
Pacific ocean perch	<u>S. alutus</u>
Redbanded rockfish	<u>S. babcocki</u>
Shortraker rockfish	<u>S. borealis</u>
Silvergray rockfish	<u>S. brevispinis</u>
Darkblotched rockfish	<u>S. crameri</u>
Splitnose rockfish	<u>S. diploproa</u>
Greenstriped rockfish	<u>S. elongatus</u>
Widow rockfish	<u>S. entomelas</u>
Yellowtail rockfish	<u>S. flavidus</u>
Rosethorn rockfish	<u>S. helvomaculatus</u>
Shortbelly rockfish	<u>S. jordani</u>
Bocaccio	<u>S. paucispinis</u>
Canary rockfish	<u>S. pinniger</u>
Redstripe rockfish	<u>S. proriger</u>
Yellowmouth rockfish	<u>S. reedi</u>
Yelloweye rockfish	<u>S. ruberrimus</u>
Harlequin rockfish	<u>S. variegatus</u>
Sharpchin rockfish	<u>S. zacentrus</u>
Shortspine thornyhead	<u>Sebastolobus alascanus</u>

Flatfish

Arrowtooth flounder	<u>Atheresthes stomias</u>
Petrable sole	<u>Eopsetta jordani</u>
Rex sole	<u>Glyptocephalus zachirus</u>
Flathead sole	<u>Hippoglossoides elassodon</u>
Pacific halibut	<u>Hippoglossus stenolepis</u>
Rock sole	<u>Lepidopsetta bilineata</u>
Slender sole	<u>Lyopsetta exilis</u>
Dover sole	<u>Microstomus pacificus</u>
English sole	<u>Parophrys vetulus</u>

Roundfish

Sablefish	<u>Anoplopoma fimbria</u>
Pacific cod	<u>Gadus macrocephalus</u>
Pacific hake	<u>Merluccius productus</u>
Lingcod	<u>Ophiodon elongatus</u>
Walleye pollock	<u>Theragra chalcogramma</u>

Table 14 (cont'd)

Selachii

Black skate
Ratfish
Big skate
Longnose skate
Starry skate
Spiny dogfish

Bathyraja trachura
Hydrolagus coliei
Raja binoculata
R. rhina
R. stellata
Squalus acanthias

Other Fish

Pearly prickleback
Pacific herring
Bigmouth sculpin
Threadfin sculpin
Bigfin eelpout
Blackfin sculpin
Northern ronquil
Eulachon
Prowfish

Bryozoichthys marjorius
Clupea harengus pallasii
Hemitripterus bolini
Icelinus filamentosus
Lycodes cortezianus
Malacocottus kincaidi
Ronquilus jordani
Thaleichthys pacificus
Zaprora silenus

Invertebrates

Squid

Octopus
Prawn

Ommastrephes bartrami
Rossia pacifica
Octopus sp.
Pandalus platyceros

C





Fig. 1. Haul locations in Queen Charlotte Sound, F/V OCEAN SELECTOR, July 12 to 30, 1989







Fig. 2. Haul locations off the west coast of Vancouver Island, F/V OCEAN SELECTOR, July 12 to 30, 1989.






Fig. 3. Haul locations off the west coast of the Queen Charlotte Islands, F/V OCEAN SELECTOR, July 12 to 30, 1989. The 400 m contour is estimated only.





Fig. 4. Schematic illustration of relative catch rates of rockfishes off Langara, F/V OCEAN SELECTOR, July 12 to 30, 1989.







Fig. 5. Schematic illustration of relative catch rates of rockfishes off Rennell Sound, F/V OCEAN SELECTOR, July 12 to 30, 1989.







Fig. 6. Schematic illustration of relative catch rates of rockfishes off Anthony Island, F/V OCEAN SELECTOR, July 12 to 30, 1989.







Fig. 7. Schematic illustration of relative catch rates of rockfishes in Moresby Gully, F/V OCEAN SELECTOR, July 12 to 30, 1989.







Fig. 8. Schematic illustration of relative catch rates of rockfishes in Mitchell's Gully, F/V OCEAN SELECTOR, July 12 to 30, 1989.







Fig. 9. Schematic illustration of relative catch rates of rockfishes in Goose Island Gully, F/V OCEAN SELECTOR, July 12 to 30, 1989.






Fig. 10. Schematic illustration of relative catch rates of rockfishes off Quatsino, F/V OCEAN SELECTOR, July 12 to 30, 1989.






Fig. 11. Schematic illustration of relative catch rates of rockfishes off southwest Vancouver Island, F/V OCEAN SELECTOR, July 12 to 30, 1989.



Appendix Table 1. Vessel specifications for the F/V OCEAN
SELECTOR.

Skipper: Dave Clattenburg
Overall Length: 47.9 m (157 ft)
Gross Tons: 452
Horsepower: 850
Capacity: 204,117 kg (450,000 lb)
Total Crew: 6
Fish Storage: Ice, Slush

ELECTRONIC AIDS

Sounder: Epsco Chromascope
Radio: Multiple
Radar: Furuno
Autopilot: Wagner
Loran: Raynav 750
Plotter: Epsco C-Plot 2, JRC NWU-52R
Net Sounder: Furuno

TRAWL GEAR

Double gear and two single winches
Cable per spool: 1,828 m (1,000 fm)
Diameter: 2.86 cm (1.125 in)
Net Reel: standard hydraulic

Appendix Table 2. Bridge log information from the slope rockfish sampling cruise aboard the F/V OCEAN SELECTOR, July 12-30, 1989.

Haul No.	1	2	3	4
Date	JUL 14	JUL 14	JUL 14	JUL 14
Area (Major, Minor)	5B,08	5B,08	5B,08	5B,08
Start Time (PDT)	7:24	10:30	12:00	21:03
Duration (Min)	41	25	116	79
Start N. Lat. (Deg)	51	51	51	51
(Min)	44.2	48.5	48.5	41.8
W. Long (Deg)	130	130	130	130
(Min)	46.8	42.6	42.6	43.3
Start LORAN-C X	12787.2	12780.5	12780.5	12814.9
Y	30174.1	30191.4	30191.4	30166.0
Finish N. Lat. (Deg)	51	51	51	51
(Min)	42.4	47.9	42.7	47.1
W. Long (Deg)	130	130	130	130
(Min)	44.0	43.6	43.9	43.7
Finish LORAN-C X	12808.7	12779.7	12807.4	12783.8
Y	30168.8	30188.8	30169.0	30185.9
Haul Distance (Km)	4.6	1.7	10.9	9.8
(N Mi)	2.5	0.9	5.9	5.3
Direction (Deg.True)	112	241	246	16
Bottom Depth (m)	275- 293	293- 329	302- 304	333- 335
(Fm)	150- 160	160- 180	165- 166	182- 183
Modal Depth (m)	275	302	320	338
Net/Gear Depth (m)	..- ..	..- ..	..- ..	..- ..
(Fm)	..- ..	..- ..	..- ..	..- ..
Modal Net Depth (m)	..	..	..	..
Footrope-Bottom (m)	0- 0	0- 0	0- 0	0- 0
Distance (Fm)	0- 0	0- 0	0- 0	0- 0
Modal Distance (m)	0	0	0	0
Gear Type	BT	BT	BT	BT
Tide	FLOOD	SLACK	EBB	FLOOD
Total Catch (Kg)	9072	11340	43091	4536
Remarks	SNAG-USE	SNAG-USE	USABLE	TORN-USE

Appendix Table 2 (cont'd)

Haul No.	1	2	3	4
Date	JUL 14	JUL 14	JUL 14	JUL 14
Area (Major, Minor)	5B,08	5B,08	5B,08	5B,08
Arrowtooth flounder	27	..	..	..
Dover sole	80	..	..	59
English sole	..	..	..	..
Flathead sole	..	..	..	..
Pacific halibut	..	..	..	..
Rex sole	..	..	..	..
Slender sole	..	..	..	..
Other Flatfish	..	..	..	..
<u>Sebastes aleutianus</u>	..	..	..	377
<u>S. alutus</u>	7608	10123	42716	2992
<u>S. babcocki</u>	53	119	..	236
<u>S. borealis</u>	293	..	..	483
<u>S. brevispinis</u>	..	..	..	..
<u>S. cramerii</u>	..	..	..	24
<u>S. diploproa</u>	..	..	..	..
<u>S. elongatus</u>	..	..	..	..
<u>S. entomelas</u>	..	..	..	..
<u>S. flavidus</u>	..	..	..	..
<u>S. helvomaculatus</u>	..	..	..	..
<u>S. paucispinis</u>	..	237	..	..
<u>S. pinniger</u>	..	..	..	..
<u>S. proriger</u>	..	..	..	..
<u>S. reedi</u>	..	178	375	..
<u>S. variegatus</u>	..	..	..	..
<u>S. zacentrus</u>	559	..	..	..
<u>Sebastolobus alascanus</u>	452	..	..	365
Other Rockfish	..	..	..	..
Eulachon	..	..	..	..
Lingcod	..	683	..	..
Pacific cod	..	..	..	..
Pacific hake	..	..	..	..
Prowfish	..	..	..	..
Sablefish	..	..	..	..
Walleye pollock	..	..	..	..
Other Roundfish	..	..	..	..
Longnose skate	..	..	..	..
Ratfish	..	..	..	..
Spiny dogfish	..	..	..	..
Starry skate	..	..	..	..
Other Selachii	..	..	..	..
Squid	..	..	..	..
Other Invertebrates	..	..	..	..
Total Catch (Kg)	9072	11340	43091	4536

Appendix Table 2 (cont'd)

Haul No.	5	6	7	8
Date	JUL 15	JUL 15	JUL 15	JUL 15
Area (Major, Minor)	5B,08	5B,08	5B,08	5C,02
Start Time (PDT)	9:20	13:05	17:50	21:45
Duration (Min)	127	127	120	80
Start N. Lat. (Deg)	51	52	52	52
(Min)	59.7	4.7	0.6	17.6
W. Long (Deg)	130	130	130	130
(Min)	38.8	32.6	13.7	3.2
Start LORAN-C X	12732.2	13049.6	12835.6	12781.3
Y	30232.8	30028.1	30249.7	30314.9
Finish N. Lat. (Deg)	52	51	52	52
(Min)	6.2	58.9	6.8	14.2
W. Long (Deg)	130	130	130	130
(Min)	42.3	35.5	10.8	4.7
Finish LORAN-C X	12679.1	12750.7	12812.1	12795.3
Y	30253.2	30231.7	30273.2	30302.4
Haul Distance (Km)	12.8	11.3	12.0	6.5
(N Mi)	6.9	6.1	6.5	3.5
Direction (Deg.True)	57	195	11	197
Bottom Depth (m)	256- 229	278- 311	269- 278	214- 212
(Fm)	140- 125	152- 170	147- 152	117- 116
Modal Depth (m)	265	294	274	214
Net/Gear Depth (m)	..- ..	..- ..	..- ..	..- ..
(Fm)	..- ..	..- ..	..- ..	..- ..
Modal Net Depth (m)	..	..	..	..
Footrope-Bottom (m)	0- 0	0- 0	0- 0	0- 0
Distance (Fm)	0- 0	0- 0	0- 0	0- 0
Modal Distance (m)	0	0	0	0
Gear Type	BT	BT	BT	BT
Tide	FLOOD	EBB	EBB	..
Total Catch (Kg)	3629	3225	2722	6804
Remarks	USABLE	USABLE	USABLE	USABLE

Appendix Table 2 (cont'd)

Haul No.	5	6	7	8
Date	JUL 15	JUL 15	JUL 15	JUL 15
Area (Major, Minor)	5B,08	5B,08	5B,08	5C,02
Arrowtooth flounder	83	50	49	496
Dover sole	..	..	29	..
English sole	12	..	..	..
Flathead sole	..	..	..	..
Pacific halibut	..	..	..	..
Rex sole	83	10	..	20
Slender sole	..	..	..	..
Other Flatfish	..	..	..	..
<u>Sebastes aleutianus</u>	..	..	..	..
<u>S. alutus</u>	3132	2547	1785	3729
<u>S. babcocki</u>	71	10	254	20
<u>S. borealis</u>	..	134	..	..
<u>S. brevispinis</u>	12	21	10	1805
<u>S. crameri</u>	..	72	..	..
<u>S. diploproa</u>	..	247	459	..
<u>S. elongatus</u>	..	..	..	..
<u>S. entomelas</u>	..	..	..	79
<u>S. flavidus</u>	..	..	..	456
<u>S. helvomaculatus</u>	..	..	..	..
<u>S. paucispinis</u>	..	..	..	..
<u>S. pinniger</u>	..	..	..	159
<u>S. proriger</u>	..	..	..	..
<u>S. reedi</u>	12	21	..	..
<u>S. variegatus</u>	..	..	..	..
<u>S. zacentrus</u>	24	..	..	..
<u>Sebastolobus alascanus</u>	47	103	78	..
Other Rockfish	..	..	..	..
Eulachon	..	..	..	..
Lingcod	..	..	..	..
Pacific cod	35	..	..	..
Pacific hake	..	..	..	..
Prowfish	..	..	..	..
Sablefish	71	10	29	40
Walleye pollock	47	..	29	..
Other Roundfish	..	..	..	..
Longnose skate	..	..	..	..
Ratfish	..	..	..	..
Spiny dogfish	..	..	..	..
Starry skate	..	..	..	..
Other Selachii	..	..	..	..
Squid	..	..	..	..
Other Invertebrates	..	..	..	..
Total Catch (Kg)	3629	3225	2722	6804

Appendix Table 2 (cont'd)

Haul No.	9	10	11	12
Date	JUL 16	JUL 16	JUL 16	JUL 17
Area (Major, Minor)	5B,08	5B,08	5B,08	3D,27
Start Time (PDT)	15:05	18:52	23:36	14:05
Duration (Min)	132	140	189	135
Start N. Lat. (Deg)	51	51	51	50
(Min)	46.7	41.5	39.9	29.8
W. Long (Deg)	129	130	129	128
(Min)	59.5	13.8	47.7	32.8
Start LORAN-C X	12978.4	12942.7	13069.6	13735.0
Y	30206.7	30179.7	30186.8	29893.9
Finish N. Lat. (Deg)	51	51	51	50
(Min)	43.0	45.0	40.9	24.2
W. Long (Deg)	130	130	129	128
(Min)	9.7	2.4	45.6	27.1
Finish LORAN-C X	12952.7	12974.6	13074.1	13781.5
Y	30187.4	30198.7	30191.9	29864.5
Haul Distance (Km)	13.5	14.6	15.2	12.4
(N Mi)	7.3	7.9	8.2	6.7
Direction (Deg.True)	242	68	56	166
Bottom Depth (m)	247- 243	225- 280	267- 265	201- 216
(Fm)	135- 133	123- 153	146- 145	110- 118
Modal Depth (m)	245	260	267	210
Net/Gear Depth (m)	..- ..	..- ..	..- ..	..- ..
(Fm)	..- ..	..- ..	..- ..	..- ..
Modal Net Depth (m)	..	..	..	..
Footrope-Bottom (m)	0- 0	0- 0	0- 0	0- 0
Distance (Fm)	0- 0	0- 0	0- 0	0- 0
Modal Distance (m)	0	0	0	0
Gear Type	BT	BT	BT	BT
Tide	SLACK	EBB	FLOOD	SLACK
Total Catch (Kg)	2948	3402	2268	6804
Remarks	USABLE	SNAG-USE	USABLE	USABLE

Appendix Table 2 (cont'd)

Haul No.	9	10	11	12
Date	JUL 16	JUL 16	JUL 16	JUL 17
Area (Major, Minor)	5B,08	5B,08	5B,08	3D,27
Arrowtooth flounder	133	11	63	151
Dover sole	19	33	32	38
English sole	..	..	..	..
Flathead sole	..	..	..	..
Pacific halibut	..	..	..	..
Rex sole	19	22	..	19
Slender sole	..	..	..	..
Other Flatfish	..	..	..	..
<u>Sebastes aleutianus</u>	..	..	25	..
<u>S. alutus</u>	2238	2128	1210	2530
<u>S. babcocki</u>	47	11	13	..
<u>S. borealis</u>	..	..	..	..
<u>S. brevispinis</u>	38	121	..	265
<u>S. crameri</u>	..	..	..	..
<u>S. diploproa</u>	..	219	38	..
<u>S. elongatus</u>	..	..	..	..
<u>S. entomelas</u>	..	..	..	..
<u>S. flavidus</u>	19	..	..	..
<u>S. helvomaculatus</u>	38	..	..	..
<u>S. paucispinis</u>	..	..	..	..
<u>S. pinniger</u>	..	..	..	..
<u>S. proriger</u>	..	11	88	794
<u>S. reedi</u>	47	450	69	813
<u>S. variegatus</u>	..	..	..	..
<u>S. zacentrus</u>	..	99	13	1058
<u>Sebastolobus alascanus</u>	95	77	6	76
Other Rockfish	..	..	..	..
Eulachon	..	..	..	..
Lingcod	..	..	..	662
Pacific cod	..	..	..	19
Pacific hake	47	33	271	227
Prowfish	..	..	..	..
Sablefish	66	132	372	95
Walleye pollock	133	44	44	..
Other Roundfish	..	..	..	..
Longnose skate	..	..	..	..
Ratfish	..	..	..	..
Spiny dogfish	..	..	..	..
Starry skate	9	11	..	57
Other Selachii	..	..	..	..
Squid	..	..	24	..
Other Invertebrates	..	..	..	..
Total Catch (Kg)	2948	3402	2268	6804

Appendix Table 2 (cont'd)

Haul No.		13	14	15	16
Date		JUL 17	JUL 18	JUL 18	JUL 18
Area (Major, Minor)		3D,27	3D,27	3D,27	3C,25
Start Time (PDT)		17:30	6:05	9:55	22:00
Duration (Min)		135	150	43	60
Start N. Lat. (Deg)		50	50	50	49
	(Min)	21.3	29.4	18.8	26.3
W. Long (Deg)		128	128	128	127
	(Min)	20.0	32.6	17.1	14.9
Start LORAN-C	X	13824.9	13737.3	13847.0	14264.0
	Y	29849.5	29891.6	29835.2	29500.8
Finish N. Lat. (Deg)		50	50	50	49
	(Min)	24.9	23.6	21.0	23.2
W. Long (Deg)		128	128	128	127
	(Min)	28.3	25.2	17.7	14.6
Finish LORAN-C	X	13773.4	13792.8	13837.2	14270.6
	Y	29868.2	29861.5	29847.3	29480.5
Haul Distance (Km)		11.8	13.9	4.1	5.8
	(N Mi)	6.4	7.5	2.2	3.1
Direction (Deg.True)		242	158	307	165
Bottom Depth (m)		254- 238	205- 208	256- 199	271- 294
	(Fm)	139- 130	112- 114	140- 109	148- 161
Modal Depth (m)		247	210	256	289
Net/Gear Depth (m)		..- ..	..- ..	..- ..	..- ..
	(Fm)	..- ..	..- ..	..- ..	..- ..
Modal Net Depth (m)		..	..	..	..
Footrope-Bottom (m)		0- 0	0- 0	0- 0	0- 0
Distance (Fm)		0- 0	0- 0	0- 0	0- 0
Modal Distance (m)		0	0	0	0
Gear Type		BT	BT	BT	BT
Tide		EBB	FLOOD	FLOOD	EBB
Total Catch (Kg)		6849	6804	3402	2722
Remarks		USABLE	USABLE	USABLE	USABLE

Appendix Table 2 (cont'd)

Haul No.	13	14	15	16
Date	JUL 17	JUL 18	JUL 18	JUL 18
Area (Major, Minor)	3D,27	3D,27	3D,27	3C,25
Arrowtooth flounder	44	101	61	448
Dover sole	..	20	..	142
English sole	..	..	..	..
Flathead sole	..	..	..	..
Pacific halibut	..	..	..	..
Rex sole	22	..	..	39
Slender sole	..	..	..	..
Other Flatfish	..	..	..	..
<u>Sebastes aleutianus</u>	..	..	..	..
<u>S. alutus</u>	3228	3744	1879	849
<u>S. babcocki</u>	..	20	8	16
<u>S. borealis</u>	..	..	..	..
<u>S. brevispinis</u>	..	221	..	..
<u>S. crameri</u>	436	..	23	8
<u>S. diploproa</u>	2115	..	1280	409
<u>S. elongatus</u>	..	20	..	..
<u>S. entomelas</u>	..	..	..	..
<u>S. flavidus</u>	..	..	..	..
<u>S. helvomaculatus</u>	22	20	8	..
<u>S. paucispinis</u>	..	322	..	..
<u>S. pinniger</u>	..	..	..	..
<u>S. proriger</u>	..	322	..	..
<u>S. reedi</u>	..	263	..	..
<u>S. variegatus</u>	..	..	..	..
<u>S. zacentrus</u>	109	966	30	8
<u>Sebastolobus alascanus</u>	218	20	45	102
Other Rockfish	..	..	..	..
Eulachon	..	..	..	..
Lingcod	..	181	..	..
Pacific cod	..	..	..	..
Pacific hake	44	..	..	189
Prowfish	..	..	..	..
Sablefish	22	262	68	488
Walleye pollock	44	20	..	24
Other Roundfish	..	..	..	..
Longnose skate	..	262	..	..
Ratfish	..	40	..	..
Spiny dogfish	..	..	..	..
Starry skate	..	..	..	..
Other Selachii	..	..	..	..
Squid	22	..	..	..
Other Invertebrates	..	..	..	..
Total Catch (Kg)	6849	6804	3402	2722

Appendix Table 2 (cont'd)

Haul No.		17	18	19	20
Date		JUL 19	JUL 19	JUL 19	JUL 19
Area (Major, Minor)		3C,25	3C,23	3C,23	3C,21
Start Time (PDT)		1:40	13:17	16:00	20:25
Duration (Min)		100	120	155	95
Start N. Lat. (Deg)		49	48	48	48
	(Min)	35.3	23.7	24.7	20.1
W. Long (Deg)		127	126	125	125
	(Min)	16.4	3.5	58.3	51.4
Start LORAN-C	X	14241.2	14626.0	14645.1	14672.7
	Y	29559.6	28992.3	28990.0	28945.6
Finish N. Lat. (Deg)		49	48	48	48
	(Min)	30.4	23.7	27.1	16.0
W. Long (Deg)		127	126	126	125
	(Min)	13.5	1.7	7.3	46.7
Finish LORAN-C	X	14263.0	14632.8	14609.8	14691.4
	Y	29526.1	28989.3	29022.8	28907.8
Haul Distance (Km)		9.8	10.8	12.0	9.6
	(N Mi)	5.3	5.9	6.5	5.2
Direction (Deg.True)		144	239	257	185
Bottom Depth (m)		159- 168	269- 229	205- 243	187- 190
	(Fm)	87- 92	147- 125	112- 133	102- 104
Modal Depth (m)		165	283	229	183
Net/Gear Depth (m)		108- 110	..- ..	..- ..	..- ..
	(Fm)	59- 60	..- ..	..- ..	..- ..
Modal Net Depth (m)		110	..	..	..
Footrope-Bottom (m)		18- 18	0- 0	0- 0	0- 0
Distance (Fm)		10- 10	0- 0	0- 0	0- 0
Modal Distance (m)		18	0	0	0
Gear Type		MW	BT	BT	BT
Tide		FLOOD	FLOOD	EBB	SLACK
Total Catch (Kg)		6804	3629	2722	2268
Remarks		USABLE	USABLE	USABLE	USABLE

Appendix Table 2 (cont'd)

Haul No.	17	18	19	20
Date	JUL 19	JUL 19	JUL 19	JUL 19
Area (Major, Minor)	3C,25	3C,23	3C,23	3C,21
Arrowtooth flounder	..	113	72	154
Dover sole	..	21	57	136
English sole	..	..	..	87
Flathead sole	..	..	..	..
Pacific halibut	..	..	..	..
Rex sole	..	62	57	118
Slender sole	..	..	..	19
Other Flatfish	..	..	..	..
<u>Sebastes aleutianus</u>	..	..	..	..
<u>S. alutus</u>	..	1996	1416	440
<u>S. babcocki</u>	..	21	79	..
<u>S. borealis</u>	..	..	..	..
<u>S. brevispinis</u>	..	..	..	31
<u>S. crameri</u>	..	185	..	..
<u>S. diploproa</u>	..	779	352	..
<u>S. elongatus</u>	..	..	29	167
<u>S. entomelas</u>	4871	..	..	6
<u>S. flavidus</u>	1933	..	..	..
<u>S. helvomaculatus</u>	..	21	36	68
<u>S. paucispinis</u>	..	..	..	..
<u>S. pinniger</u>	..	..	..	..
<u>S. proriger</u>	..	..	..	192
<u>S. reedi</u>	..	..	..	647
<u>S. variegatus</u>	..	..	..	..
<u>S. zacentrus</u>	..	10	136	74
<u>Sebastolobus alascanus</u>	..	82	158	..
Other Rockfish	..	..	..	..
Eulachon	..	..	..	..
Lingcod	..	..	..	..
Pacific cod	..	..	29	49
Pacific hake	..	41	..	12
Prowfish	..	..	..	..
Sablefish	..	236	215	31
Walleye pollock	..	21	36	..
Other Roundfish	..	..	..	..
Longnose skate	..	..	..	..
Ratfish	..	41	..	31
Spiny dogfish	..	..	50	6
Starry skate	..	..	..	..
Other Selachii	..	..	..	..
Squid	..	..	..	..
Other Invertebrates	..	..	..	..
Total Catch (Kg)	6804	3629	2722	2268

Appendix Table 2 (cont'd)

Haul No.		21	22	23	24
Date		JUL 19	JUL 20	JUL 26	JUL 26
Area (Major, Minor)		3C,21	3C,21	5E,35	5E,35
Start Time (PDT)		22:25	1:35	7:40	10:15
Duration (Min)		165	95	60	125
Start N. Lat. (Deg)		48	48	54	54
	(Min)	15.5	20.1	5.3	9.7
W. Long (Deg)		125	125	133	133
	(Min)	46.6	51.2	42.9	45.3
Start LORAN-C	X	14692.1	14673.7	11414.5	11387.4
	Y	28904.5	28945.1	30466.7	30474.2
Finish N. Lat. (Deg)		48	48	54	54
	(Min)	20.1	17.9	2.6	16.1
W. Long (Deg)		125	125	133	133
	(Min)	51.1	50.9	40.7	49.7
Finish LORAN-C	X	14673.9	14675.5	11432.1	11348.8
	Y	28945.0	28929.3	30462.3	30484.4
Haul Distance (Km)		10.2	4.1	5.6	12.8
	(N Mi)	5.5	2.2	3.0	6.9
Direction (Deg.True)		278	214	169	297
Bottom Depth (m)		201- 183	198- 201	318- 327	263- 260
	(Fm)	110- 100	108- 110	174- 179	144- 142
Modal Depth (m)		192	199	338	262
Net/Gear Depth (m)		..- ..	..- ..	..- ..	..- ..
	(Fm)	..- ..	..- ..	..- ..	..- ..
Modal Net Depth (m)		..	..	..	..
Footrope-Bottom (m)		0- 0	0- 0	0- 0	0- 0
Distance (Fm)		0- 0	0- 0	0- 0	0- 0
Modal Distance (m)		0	0	0	0
Gear Type		BT	BT	BT	BT
Tide		EBB	EBB	SLACK	EBB
Total Catch (Kg)		9072	2268	9072	6804
Remarks		USABLE	USABLE	USABLE	USABLE

Appendix Table 2 (cont'd)

Haul No.	21	22	23	24
Date	JUL 19	JUL 20	JUL 26	JUL 26
Area (Major, Minor)	3C,21	3C,21	5E,35	5E,35
Arrowtooth flounder	536	113	132	..
Dover sole	382	..	..	..
English sole	38	..	..	..
Flathead sole	..	..	..	..
Pacific halibut	..	..	..	45
Rex sole	153	..	..	..
Slender sole	..	..	..	..
Other Flatfish	..	..	..	..
<u>Sebastes aleutianus</u>	..	..	7115	..
<u>S. alutus</u>	4249	680	1746	6088
<u>S. babcocki</u>	77	..	..	80
<u>S. borealis</u>	..	..	..	..
<u>S. brevispinis</u>	..	..	..	45
<u>S. crameri</u>	..	..	..	..
<u>S. diploproa</u>	..	454	..	..
<u>S. elongatus</u>	421	..	..	..
<u>S. entomelas</u>	..	..	..	..
<u>S. flavidus</u>	..	..	..	..
<u>S. helvomaculatus</u>	..	..	..	..
<u>S. paucispinis</u>	..	..	..	..
<u>S. pinniger</u>	..	..	..	..
<u>S. proriger</u>	115	..	..	..
<u>S. reedi</u>	1340	908	..	..
<u>S. variegatus</u>	..	..	..	..
<u>S. zacentrus</u>	421	..	..	381
<u>Sebastolobus alascanus</u>	268	..	79	120
Other Rockfish	..	..	..	..
Eulachon	..	..	..	..
Lingcod	..	..	..	45
Pacific cod	..	..	..	..
Pacific hake	230	..	..	..
Prowfish	..	..	..	..
Sablefish	689	113	..	..
Walleye pollock	..	..	..	..
Other Roundfish	..	..	..	..
Longnose skate	..	..	..	..
Ratfish	..	..	..	..
Spiny dogfish	..	..	..	..
Starry skate	..	..	..	..
Other Selachii	..	..	..	..
Squid	..	..	..	..
Other Invertebrates	..	..	..	..
Total Catch (Kg)	9072	2268	9072	6804

Appendix Table 2 (cont'd)

Haul No.	25	26	27	28
Date	JUL 26	JUL 26	JUL 26	JUL 27
Area (Major, Minor)	5E,35	5E,35	5E,35	5E,35
Start Time (PDT)	12:47	15:50	20:15	8:27
Duration (Min)	110	180	110	109
Start N. Lat. (Deg)	54	54	54	53
(Min)	17.2	22.2	9.3	24.3
W. Long (Deg)	133	133	133	133
(Min)	49.7	51.6	45.8	9.6
Start LORAN-C X	11342.8	11315.1	11389.2	11712.8
Y	30486.5	30495.0	30473.2	30394.6
Finish N. Lat. (Deg)	54	54	54	53
(Min)	22.7	12.9	13.9	19.5
W. Long (Deg)	133	133	133	133
(Min)	51.0	48.9	50.2	4.3
Finish LORAN-C X	11312.8	11366.8	11360.3	11753.9
Y	30496.3	30478.7	30479.0	30385.0
Haul Distance (Km)	10.4	17.4	9.8	10.7
(N Mi)	5.6	9.4	5.3	5.8
Direction (Deg.True)	357	202	313	171
Bottom Depth (m)	238- 271	252- 263	302- 338	219- 227
(Fm)	130- 148	138- 144	165- 185	120- 124
Modal Depth (m)	238	256	311	223
Net/Gear Depth (m)	..- ..	..- ..	..- ..	..- ..
(Fm)	..- ..	..- ..	..- ..	..- ..
Modal Net Depth (m)	..	..	..	..
Footrope-Bottom (m)	0- 0	0- 0	0- 0	0- 0
Distance (Fm)	0- 0	0- 0	0- 0	0- 0
Modal Distance (m)	0	0	0	0
Gear Type	BT	BT	BT	BT
Tide	EBB	FLOOD	SLACK	FLOOD
Total Catch (Kg)	2722	7257	4990	14515
Remarks	SNAG-USE	SNAG-USE	SNAG-USE	USABLE

Appendix Table 2 (cont'd)

Haul No.	25	26	27	28
Date	JUL 26	JUL 26	JUL 26	JUL 27
Area (Major, Minor)	5E,35	5E,35	5E,35	5E,35
Arrowtooth flounder	40	23	80	84
Dover sole	..	..	133	..
English sole	..	..	..	..
Flathead sole	..	..	..	..
Pacific halibut	..	..	..	..
Rex sole	..	..	..	..
Slender sole	..	..	..	..
Other Flatfish	..	..	..	..
<u>Sebastes aleutianus</u>	..	68	240	..
<u>S. alutus</u>	1017	5854	3351	755
<u>S. babcocki</u>	..	90	27	..
<u>S. borealis</u>	..	..	506	..
<u>S. brevispinis</u>	202	113	..	336
<u>S. crameri</u>	..	..	27	..
<u>S. diploproa</u>	..	..	..	..
<u>S. elongatus</u>	..	..	..	..
<u>S. entomelas</u>	..	23	..	252
<u>S. flavidus</u>	..	..	..	..
<u>S. helvomaculatus</u>	..	..	27	84
<u>S. paucispinis</u>	..	..	..	..
<u>S. pinniger</u>	..	..	..	..
<u>S. proriger</u>	16	23	..	2727
<u>S. reedi</u>	8	68	..	10235
<u>S. variegatus</u>	145	..	..	..
<u>S. zacentrus</u>	1197	972	386	..
<u>Sebastolobus alascanus</u>	40	23	200	42
Other Rockfish	..	..	..	..
Eulachon	..	..	..	..
Lingcod	..	..	..	..
Pacific cod	57	..	..	..
Pacific hake	..	..	..	..
Prowfish	..	..	..	..
Sablefish	..	..	..	..
Walleye pollock	..	..	..	..
Other Roundfish	..	..	13	..
Longnose skate	..	..	..	..
Ratfish	..	..	..	..
Spiny dogfish	..	..	..	..
Starry skate	..	..	..	..
Other Selachii	..	..	..	..
Squid	..	..	..	..
Other Invertebrates	..	..	..	..
Total Catch (Kg)	2722	7257	4990	14515

Appendix Table 2 (cont'd)

Haul No.		29	30	31	32
Date		JUL 27	JUL 27	JUL 27	JUL 28
Area (Major, Minor)		5E,31	5E,31	5E,31	5E,34
Start Time (PDT)		12:37	16:05	19:32	10:45
Duration (Min)		167	163	218	70
Start N. Lat. (Deg)		53	53	53	52
	(Min)	18.2	25.7	16.9	2.0
W. Long (Deg)		133	133	133	131
	(Min)	2.9	10.7	2.7	20.3
Start LORAN-C	X	11765.1	11701.8	11773.8	12553.4
	Y	30383.6	30397.4	30380.6	30219.9
Finish N. Lat. (Deg)		53	53	53	51
	(Min)	25.4	17.9	26.1	59.1
W. Long (Deg)		133	133	133	131
	(Min)	10.5	5.4	11.0	16.3
Finish LORAN-C	X	11704.1	11761.8	11698.7	12585.5
	Y	30396.8	30381.6	30398.1	30212.2
Haul Distance (Km)		15.7	15.6	19.4	7.0
	(N Mi)	8.5	8.4	10.5	3.8
Direction (Deg.True)		245	170	273	142
Bottom Depth (m)		280- 276	300- 280	254- 265	241- 293
	(Fm)	153- 151	164- 153	139- 145	132- 160
Modal Depth (m)		278	309	260	256
Net/Gear Depth (m)		..- ..	..- ..	..- ..	..- ..
	(Fm)	..- ..	..- ..	..- ..	..- ..
Modal Net Depth (m)		..	..	..	..
Footrope-Bottom (m)		0- 0	0- 0	0- 0	0- 0
Distance (Fm)		0- 0	0- 0	0- 0	0- 0
Modal Distance (m)		0	0	0	0
Gear Type		BT	BT	BT	BT
Tide		EBB	EBB	FLOOD	EBB
Total Catch (Kg)		5897	12247	6804	6804
Remarks		USABLE	USABLE	USABLE	USABLE

Appendix Table 2 (cont'd)

Haul No.	29	30	31	32
Date	JUL 27	JUL 27	JUL 27	JUL 28
Area (Major, Minor)	5E,31	5E,31	5E,31	5E,34
Arrowtooth flounder	..	..	..	..
Dover sole	..	..	..	..
English sole	..	..	..	..
Flathead sole	..	..	..	..
Pacific halibut	..	..	..	..
Rex sole	..	..	..	..
Slender sole	..	..	..	..
Other Flatfish	..	..	..	..
<u>Sebastes aleutianus</u>	..	233	136	..
<u>S. alutus</u>	4415	11432	5784	5630
<u>S. babcocki</u>	16	78	..	78
<u>S. borealis</u>	..	..	136	..
<u>S. brevispinis</u>	65	..	340	98
<u>S. crameri</u>	..	..	..	..
<u>S. diploproa</u>	..	..	..	..
<u>S. elongatus</u>	..	..	..	..
<u>S. entomelas</u>	..	..	..	20
<u>S. flavidus</u>	..	..	..	..
<u>S. helvomaculatus</u>	98	..	68	20
<u>S. paucispinis</u>	81	..	..	..
<u>S. pinniger</u>	..	..	..	..
<u>S. proriger</u>	98	..	..	156
<u>S. reedi</u>	782	..	..	156
<u>S. variegatus</u>	..	..	..	..
<u>S. zacentrus</u>	..	..	..	411
<u>Sebastolobus alascanus</u>	114	504	..	117
Other Rockfish	..	..	..	..
Eulachon	..	..	..	..
Lingcod	..	..	..	..
Pacific cod	..	..	..	..
Pacific hake	..	..	..	..
Prowfish	81	..	..	..
Sablefish	147	..	340	98
Walleye pollock	..	..	..	20
Other Roundfish	..	..	..	..
Longnose skate	..	..	..	..
Ratfish	..	..	..	..
Spiny dogfish	..	..	..	..
Starry skate	..	..	..	..
Other Selachii	..	..	..	..
Squid	..	..	..	..
Other Invertebrates	..	..	..	..
Total Catch (Kg)	5897	12247	6804	6804

Appendix Table 2 (cont'd)

Haul No.	33	34	35	36
Date	JUL 28	JUL 28	JUL 28	JUL 28
Area (Major, Minor)	5E,34	5C,02	5C,02	5C,02
Start Time (PDT)	12:28	19:02	20:15	23:07
Duration (Min)	96	48	145	80
Start N. Lat. (Deg)	51	52	52	52
(Min)	59.3	17.7	20.9	20.8
W. Long (Deg)	131	130	130	130
(Min)	16.2	23.2	23.7	22.6
Start LORAN-C X	12584.8	12690.9	12668.8	12674.2
Y	30212.9	30302.6	30312.8	30313.2
Finish N. Lat. (Deg)	52	52	52	52
(Min)	2.9	20.7	20.4	25.1
W. Long (Deg)	131	130	130	130
(Min)	21.6	23.6	23.0	22.4
Finish LORAN-C X	12543.2	12670.5	12674.0	12648.1
Y	30222.3	30312.2	30311.6	30327.2
Haul Distance (Km)	9.1	5.6	13.0	8.0
(N Mi)	4.9	3.0	6.6	4.3
Direction (Deg.True)	334	24	150	353
Bottom Depth (m)	327- 371	227- 245	248- 245	290- 274
(Fm)	179- 203	124- 134	136- 134	159- 150
Modal Depth (m)	320	219	241	282
Net/Gear Depth (m)	..- ..	..- ..	..- ..	..- ..
(Fm)	..- ..	..- ..	..- ..	..- ..
Modal Net Depth (m)	..	..	..	..
Footrope-Bottom (m)	0- 0	0- 0	0- 0	0- 0
Distance (Fm)	0- 0	0- 0	0- 0	0- 0
Modal Distance (m)	0	0	0	0
Gear Type	BT	BT	BT	BT
Tide	FLOOD	FLOOD	FLOOD	EBB
Total Catch (Kg)	3175	2268	9979	1814
Remarks	USABLE	USABLE	USABLE	USABLE

Appendix Table 2 (cont'd)

Haul No.	33	34	35	36
Date	JUL 28	JUL 28	JUL 28	JUL 28
Area (Major, Minor)	5E,34	5C,02	5C,02	5C,02
Arrowtooth flounder	29	416	70	107
Dover sole	..	7	70	78
English sole	..	..	..	..
Flathead sole	..	..	..	..
Pacific halibut	..	..	..	..
Rex sole	..	7	..	44
Slender sole	..	..	..	..
Other Flatfish	..	..	..	..
<u>Sebastes aleutianus</u>	153	..	..	29
<u>S. alutus</u>	2485	676	8481	1032
<u>S. babcocki</u>	115	..	70	..
<u>S. borealis</u>	38	..	..	..
<u>S. brevispinis</u>	19	902	730	..
<u>S. crameri</u>	..	..	..	..
<u>S. diploproa</u>	..	..	..	..
<u>S. elongatus</u>	..	..	..	..
<u>S. entomelas</u>	..	..	..	..
<u>S. flavidus</u>	..	..	..	..
<u>S. helvomaculatus</u>	38	..	..	5
<u>S. paucispinis</u>	..	42	..	..
<u>S. pinniger</u>	..	..	..	..
<u>S. proriger</u>	29	..	..	..
<u>S. reedi</u>	..	35	..	..
<u>S. variegatus</u>	..	..	..	..
<u>S. zacentrus</u>	77	106	313	..
<u>Sebastolobus alascanus</u>	163	42	70	189
Other Rockfish	..	..	..	..
Eulachon	..	..	..	..
Lingcod	..	..	..	..
Pacific cod	..	21	35	..
Pacific hake	..	..	..	5
Prowfish	..	..	..	..
Sablefish	..	7	70	296
Walleye pollock	29	7	70	29
Other Roundfish	..	..	..	..
Longnose skate	..	..	..	..
Ratfish	..	..	..	..
Spiny dogfish	..	..	..	..
Starry skate	..	..	..	..
Other Selachii	..	..	..	..
Squid	..	..	..	..
Other Invertebrates	..	..	..	..
Total Catch (Kg)	3175	2268	9979	1814

Appendix Table 2 (cont'd)

Haul No.	37	38	39	40
Date	JUL 29	JUL 29	JUL 29	JUL 29
Area (Major, Minor)	5B,08	5B,08	5B,08	5B,08
Start Time (PDT)	8:56	12:04	16:00	19:30
Duration (Min)	131	148	135	190
Start N. Lat. (Deg)	51	51	51	51
(Min)	25.3	27.4	17.4	22.1
W. Long (Deg)	129	128	129	129
(Min)	1.4	39.5	6.6	22.4
Start LORAN-C X	13366.9	13467.7	13378.8	13279.9
Y	30150.0	30172.5	30113.5	30126.6
Finish N. Lat. (Deg)	51	51	51	51
(Min)	26.3	23.5	17.2	25.1
W. Long (Deg)	128	128	129	129
(Min)	49.1	51.9	19.7	7.4
Finish LORAN-C X	13423.9	13423.1	13316.3	13338.2
Y	30162.1	30148.1	30106.6	30146.0
Haul Distance (Km)	14.4	16.1	15.2	18.3
(N Mi)	7.8	8.7	8.2	9.9
Direction (Deg.True)	84	154	255	89
Bottom Depth (m)	165- 177	203- 219	262- 282	143- 124
(Fm)	90- 97	111- 120	143- 154	78- 68
Modal Depth (m)	165	219	274	134
Net/Gear Depth (m)	..- ..	..- ..	..- ..	80- 80
(Fm)	..- ..	..- ..	..- ..	44- 44
Modal Net Depth (m)	..	..	..	80
Footrope-Bottom (m)	0- 0	0- 0	0- 0	33- 33
Distance (Fm)	0- 0	0- 0	0- 0	18- 18
Modal Distance (m)	0	0	0	33
Gear Type	BT	BT	BT	MW
Tide	FLOOD	SLACK	EBB	FLOOD
Total Catch (Kg)	2268	5897	1814	20412
Remarks	USABLE	USABLE	USABLE	USABLE

Appendix Table 2 (cont'd)

Haul No.	37	38	39	40
Date	JUL 29	JUL 29	JUL 29	JUL 29
Area (Major, Minor)	5B,08	5B,08	5B,08	5B,08
Arrowtooth flounder	180	702	32	..
Dover sole	84	737	19	..
English sole	..	..	..	..
Flathead sole	18	35	..	..
Pacific halibut	..	..	..	..
Rex sole	30	140	13	..
Slender sole	..	..	..	..
Other Flatfish	..	..	..	..
<u>Sebastes aleutianus</u>	..	..	45	..
<u>S. alutus</u>	1043	3248	1365	..
<u>S. babcocki</u>	295	298	45	..
<u>S. borealis</u>	..	..	13	..
<u>S. brevispinis</u>	114	105	..	..
<u>S. crameri</u>	..	..	..	..
<u>S. diploproa</u>	..	..	..	..
<u>S. elongatus</u>	..	..	..	..
<u>S. entomelas</u>	66	..	..	10983
<u>S. flavidus</u>	..	..	..	9085
<u>S. helvomaculatus</u>	..	..	..	..
<u>S. paucispinis</u>	132	..	26	115
<u>S. pinniger</u>	..	..	..	..
<u>S. proriger</u>	12	..	..	57
<u>S. reedi</u>	..	..	..	..
<u>S. variegatus</u>	..	..	..	..
<u>S. zacentrus</u>	..	..	..	..
<u>Sebastolobus alascanus</u>	72	193	51	..
Other Rockfish	..	..	..	..
Eulachon	18	..	..	..
Lingcod	..	..	..	..
Pacific cod	..	..	6	..
Pacific hake	60	..	..	115
Prowfish	..	..	..	..
Sablefish	84	439	199	..
Walleye pollock	54	..	..	57
Other Roundfish	..	..	..	..
Longnose skate	..	..	..	..
Ratfish	..	..	..	..
Spiny dogfish	6	..	..	..
Starry skate	..	..	..	..
Other Selachii	..	..	..	..
Squid	..	..	..	..
Other Invertebrates	..	..	..	..
Total Catch (Kg)	2268	5897	1814	20412

Appendix Table 3. Description of rockfish gonad maturity stages.


Code		Gonad condition
Females:	1	Immature (translucent, small)
	2	Developing (small yellow eggs, translucent or opaque)
	3	Developed (large yellow eggs, opaque)
	4	Fertilized (large orange-yellow eggs, translucent)
	5	Embryos or larvae (includes eyed eggs)
	6	Spent (large flaccid red ovaries, a few larvae may be present)
	7	Resting (moderate size, firm, red-grey ovaries)
Males:	1	Immature (translucent, string-like)
	2	Developing (swelling, brown-white)
	4	Developed (large, white; easily broken)
	5	Ripe (running sperm)
	6	Spent (flaccid, red)
	7	Resting (ribbon-like, small, brown)

Appendix Figure 1.

NET DIMENSIONS AND CHARACTERISTICS FOR BOTTOM Trawl

VESSEL OCEAN SELECTOR NET Western IIA

OBSERVATION PERIOD July 12-30, 1989



Appendix Figure 2.

NET DIMENSIONS AND CHARACTERISTICS

VESSEL NAME OCEAN SELECTOR VESSEL TYPE Trawler

July 12-30, 1989

TRAWL DOORS

Shape rectangle

Type _____

SWEETLINE /

DANDYLINE

Dimensions

2.83 m x 1.81 m

Length 91 m

Weight 1270 kg


FLOATS

Number _____

Size _____

Shape _____

Material _____



Headrope Length _____ m.

Footrope Length _____ m.

Horizontal Opening 44 m.

Vertical Opening _____ m.

"a" _____ m.

"b" _____ kg.

Length 41

WINGS

(mesh or line)

Mesh Size _____ cm.

Line Spacing _____ cm.

Line Thickness _____ cm.

Material _____

Height of Wings _____ m.

MESH SIZE

81

81

81

41

20

10 & 8

cm

Mesh Material

Polish rope trawl No. 7